Anti-Gasdermin D (N terminal) Antibody ER1901-37

Product Type:	Rabbit polyclonal IgG, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IHC-P, FC, IF-Cell
Molecular Wt:	Predicted band size: 53 kDa
Description:	Gasdermin-D, N-terminal: Promotes pyroptosis in response to microbial infection and danger signals. Produced by the cleavage of gasdermin-D by inflammatory caspases CASP1 or CASP4 in response to canonical, as well as non-canonical (such as cytosolic LPS) inflammasome activators. After cleavage, moves to the plasma membrane where it strongly binds to inner leaflet lipids, including monophosphorylated phosphatidylinositols, such as phosphatidylinositol 4-phosphate, bisphosphorylated phosphatidylinositol, such as phosphatidylinositol (4,5)-bisphosphate, as well as phosphatidylinositol (3,4,5)-bisphosphate, and more weakly to phosphatidic acid and phosphatidylserine. Homooligomerizes within the membrane and forms pores of 10 - 15 nanometers (nm) of inner diameter, possibly allowing the release of mature IL1B and triggering pyroptosis . Exhibits bactericidal activity. Gasdermin-D, N-terminal released from pyroptotic cells into the extracellular milieu rapidly binds to and kills both Gram-negative and Gram-positive bacteria, without harming neighboring mammalian cells, as it does not disrupt the plasma membrane from the outside due to lipid-binding specificity.
lmmunogen:	Recombinant protein within human Gasdermin D aa 100-300.
Positive control:	SiHa cell lysate, PC-3 cell lysate, Jurkat cell lysate, THP-1 cell lysate, human kidney tissue lysate, SiHa, rat stomach tissue, human tonsil tissue, human prostate carcinoma tissue, human esophagus tissue, mouse colon tissue, mouse intestine tissue, SiHa, PC-3M, RAW264.7, L6.
Subcellular location:	Cytosol. Secreted. Plasma membrane.
Database links:	SwissProt: P57764 Human Q9D8T2 Mouse Entrez Gene: 315084 Rat
Recommended Dilutions: WB IHC-P FC IF-Cell Storage Buffer:	1:1,000-1:2,000 1:50-1:1,000 1:1,000 1:100 1*PBS (pH7.4) 0.2% BSA 50% Glycerol Preservative: 0.05% Sodium Azide
Storage Instruction	Shipped at A° . Store at $\pm A^{\circ}$ short term (1.2 weaks). It is recommended to aligned into
	single-use upon delivery. Store at -20° long term.
Purity:	Immunogen attinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn



Images



Fig1: Western blot analysis of Gasdermin D (N terminal) on different lysates with Rabbit anti-Gasdermin D (N terminal) antibody (ER1901-37) at 1/1,000 dilution.

Lane 1: SiHa cell lysate (10 µg/Lane) Lane 2: PC-3 cell lysate (10 µg/Lane) Lane 3: Jurkat cell lysate (10 µg/Lane) Lane 4: THP-1 cell lysate (10 µg/Lane) Lane 5: Human kidney tissue lysate (20 µg/Lane)

Predicted band size: 53 kDa Observed band size: 53 kDa

Exposure time: 2 minutes;

12% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (ER1901-37) at 1/1,000 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1:300,000 dilution was used for 1 hour at room temperature.

Fig2: Western blot analysis of Gasdermin D (N terminal) on THP-1 cell lysates with Rabbit anti-Gasdermin D (N terminal) antibody (ER1901-37) at 1/1,000 dilution.

Lysates/proteins at 20 µg/Lane.

Predicted band size: 53 kDa Observed band size: 53/35 kDa

Exposure time: 25 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (ER1901-37) at 1/1,000 dilution was used in 5% NFDM/TBST at 4° C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

250 150

100

55

45

35-

25

14

Gasdermin D

N terminal

-53kDa

GAPDH

Technical:0086-571-89986345

Service mail:support@huabio.cn







Fig3: Immunohistochemical analysis of paraffin-embedded rat stomach tissue using anti-Gasdermin D (N terminal) antibody. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ER1901-37, 1/200) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Fig4: Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-Gasdermin D (N terminal) antibody. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ER1901-37, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Fig5: Immunohistochemical analysis of paraffin-embedded human prostate carcinoma tissue using anti-Gasdermin D (N terminal) antibody. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ER1901-37, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Fig6: Immunohistochemical analysis of paraffin-embedded human esophagus tissue using anti-Gasdermin D (N terminal) antibody. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ER1901-37, 1/200) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn





Fig7: Immunohistochemical analysis of paraffin-embedded Mouse colon tissue using anti-Gasdermin D (N terminal) antibody. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ER1901-37, 1/200) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Fig8: Immunohistochemical analysis of paraffin-embedded mouse intestine tissue with Rabbit anti-Gasdermin D (N terminal) antibody (ER1901-37) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ER1901-37) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Fig9: Immunocytochemistry analysis of PC-3M cells labeling Gasdermin D (N terminal) with Rabbit anti-Gasdermin D (N terminal) antibody (ER1901-37) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Gasdermin D (N terminal) antibody (ER1901-37) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at $+4^{\circ}$ C. Goat Anti-Mouse IgG H&L (iFluor 1594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

が 柴安生物 www.huabio.cn



Fig10: Immunocytochemistry analysis of RAW264.7 cells labeling Gasdermin D (N terminal) with Rabbit anti-Gasdermin D (N terminal) antibody (ER1901-37) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Gasdermin D (N terminal) antibody (ER1901-37) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at $+4^{\circ}$ C. Goat Anti-Mouse IgG H&L (iFluor 1500 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.



Fig11: Immunocytochemistry analysis of L6 cells labeling Gasdermin D (N terminal) with Rabbit anti-Gasdermin D (N terminal) antibody (ER1901-37) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Gasdermin D (N terminal) antibody (ER1901-37) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at $+4^{\circ}$ C. Goat Anti-Mouse IgG H&L (iFluor 150 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.



Fig12: Flow cytometric analysis of Gasdermin D (N terminal) was done on SiHa cells. The cells were fixed, permeabilized and stained with the primary antibody (ER1901-37, 1/50) (red). After incubation of the primary antibody at room temperature for an hour, the cells were stained with a Alexa Fluor 488-conjugated Goat anti-Rabbit IgG Secondary antibody at 1/1000 dilution for 30 minutes.Unlabelled sample was used as a control (cells without incubation with primary antibody; black).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

が 柴安生物 www.hugbio.cn

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

- 1. Sborgi L. et. al. GSDMD membrane pore formation constitutes the mechanism of pyroptotic cell death. EMBO J. 35:1766-1778(2016).
- 2. Ding J. et. al. Pore-forming activity and structural autoinhibition of the gasdermin family. Nature 535:111-116(2016).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

が 华安生物 www.huabio.cn

Technical:0086-571-89986345

45 Service mail:support@huabio.cn