
Anti-Cleaved-Factor B Bb (K260) Antibody ER60037



Species reactivity: Human, Mouse

Applications: WB

Molecular Wt: 57/85 kDa

Description: This gene encodes complement factor B, a component of the alternative pathway of

complement activation. Factor B circulates in the blood as a single chain polypeptide. Upon activation of the alternative pathway, it is cleaved by complement factor D yielding the noncatalytic chain Ba and the catalytic subunit Bb. The active subunit Bb is a serine protease which associates with C3b to form the alternative pathway C3 convertase. Bb is involved in the proliferation of preactivated B lymphocytes, while Ba inhibits their proliferation. This gene localizes to the major histocompatibility complex (MHC) class III region on chromosome 6. This cluster includes several genes involved in regulation of the immune reaction. Polymorphisms in this gene are associated with a reduced risk of age-related macular degeneration. The polyadenylation site of this gene is 421 bp from the 5'

end of the gene for complemen

Immunogen: The antiserum was produced against synthesized peptide derived from human CFAB Bb. AA

range:241-290

Positive control: Blood, Colon, Liver, Plasma,

Subcellular location: Secreted

Database links: SwissProt: P00751 Human

Recommended Dilutions:

WB 1:500-1:2000

Storage Buffer: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Storage Instruction: Store at +4℃ after thawing. Aliquot store at -20℃ or -80℃. Avoid repeated freeze / thaw

cycles.

Purity: Immunogen affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.



Service mail:support@huabio.cn



Images




Fig1: Western blot analysis of lysates from K562 cells, treated with etoposide 25uM 1h, using CFAB Bb (Cleaved-Lys260) Antibody. The lane on the right is blocked with the synthesized peptide.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".