

# Anti-Hsp70 Antibody [SA0379]

## ET1601-11



<b>Product Type:</b>	Recombinant Rabbit monoclonal IgG, primary antibodies
<b>Species reactivity:</b>	Human, Mouse, Rat
<b>Applications:</b>	WB, IF-Cell, IF-Tissue, IHC-P, FC
<b>Molecular Wt:</b>	Predicted band size: 70 kDa
<b>Clone number:</b>	SA0379

<b>Description:</b>	The 70 kilodalton heat shock proteins (Hsp70s) are a family of conserved ubiquitously expressed heat shock proteins. Proteins with similar structure exist in virtually all living organisms. The Hsp70s are an important part of the cell's machinery for protein folding, and help to protect cells from stress. When not interacting with a substrate peptide, Hsp70 is usually in an ATP bound state. Hsp70 by itself is characterized by a very weak ATPase activity, such that spontaneous hydrolysis will not occur for many minutes. As newly synthesized proteins emerge from the ribosomes, the substrate binding domain of Hsp70 recognizes sequences of hydrophobic amino acid residues, and interacts with them. This spontaneous interaction is reversible, and in the ATP bound state Hsp70 may relatively freely bind and release peptides. However, the presence of a peptide in the binding domain stimulates the ATPase activity of Hsp70, increasing its normally slow rate of ATP hydrolysis.
<b>Immunogen:</b>	Recombinant protein within Mouse Hsp70 aa 403-641 / 641.
<b>Positive control:</b>	HeLa cell lysate, A549 cell lysate, MCF7 cell lysate, HCT 116 cell lysate, mouse brain tissue lysate, mouse testis tissue lysate, rat testis tissue lysate, NIH/3T3 cell lysate, C2C12 cell lysate, HeLa, MCF-7, human breast carcinoma tissue, human liver carcinoma tissue, human breast tissue, human kidney tissue, human small intestine tissue, mouse brain tissue, mouse prostate tissue.
<b>Subcellular location:</b>	Cytoplasm
<b>Database links:</b>	SwissProt: P17879 Mouse   Q61696 Mouse   P0DMV9 Human   P0DMV8 Human   Q07439 Rat
<b>Recommended Dilutions:</b>	
<b>WB</b>	1:2,000-1:5,000
<b>IF-Cell</b>	1:100
<b>IF-Tissue</b>	1:50
<b>IHC-P</b>	1:200-1:1,000
<b>FC</b>	1:50
<b>Storage Buffer:</b>	1*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.
<b>Storage Instruction:</b>	Store at +4°C after thawing. Aliquot store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
<b>Purity:</b>	Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

## Images

**Fig1:** Western blot analysis of Hsp70 on different lysates with Rabbit anti-Hsp70 antibody (ET1601-11) at 1/2,000 dilution.



Lane 1: HeLa cell lysate  
 Lane 2: A549 cell lysate  
 Lane 3: MCF7 cell lysate  
 Lane 4: HCT 116 cell lysate  
 Lane 5: Mouse brain tissue lysate  
 Lane 6: Mouse testis tissue lysate  
 Lane 7: Rat testis tissue lysate

Lysates/proteins at 15 µg/Lane.

Predicted band size: 70 kDa


Observed band size: 70 kDa

Exposure time: Lane 1-4: 43 seconds; Lane 5-7: 2 minutes;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (ET1601-11) at 1/2,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1:50,000 dilution was used for 1 hour at room temperature.

**Fig2:** Western blot analysis of Hsp70 on different lysates with Rabbit anti-Hsp70 antibody (ET1601-11) at 1/5,000 dilution.



Lane 1: NIH/3T3 cell lysate  
 Lane 2: C2C12 cell lysate

Lysates/proteins at 20 µg/Lane.

Predicted band size: 70 kDa

Observed band size: 70 kDa

Exposure time: 5 minutes;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (ET1601-11) at 1/5,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1:50,000 dilution was used for 1 hour at room temperature.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
 HUABIO  
 www.huabio.cn

**Fig3:** Western blot analysis of Hsp70 on different lysates with Rabbit anti-Hsp70 antibody (ET1601-11) at 1/1,000 dilution.

Lane 1: A549-si NT cell lysate

Lane 2: A549-si Hsp70 cell lysate

Lysates/proteins at 10 µg/Lane.


Predicted band size: 70 kDa

Observed band size: 70 kDa

Exposure time: 49 seconds;


ECL: merk

4-20% SDS-PAGE gel.



ET1601-11 was shown to specifically react with Hsp70 in A549-si NT cells. Weakened band was observed when A549-si Hsp70 sample was tested. A549-si NT and A549-si Hsp70 samples were subjected to SDS-PAGE. Proteins were transferred to a PVDF membrane and blocked with 5% NFDM in TBST for 1 hour at room temperature. The primary antibody (ET1601-11, 1/1,000) and Loading control antibody (Rabbit anti-GAPDH, ET1601-4, 1/10,000) were used in 5% BSA at room temperature for 2 hours. Goat Anti-rabbit IgG-HRP Secondary Antibody (HA1001) at 1:100,000 dilution was used for 1 hour at room temperature.

**Fig4:** Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue with Rabbit anti-Hsp70 antibody (ET1601-11) at 1/1,000 dilution.



The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1601-11) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn




**Fig5:** Immunohistochemical analysis of paraffin-embedded human liver carcinoma tissue with Rabbit anti-Hsp70 antibody (ET1601-11) at 1/1,000 dilution.


The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1601-11) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




**Fig6:** Immunohistochemical analysis of paraffin-embedded human breast tissue using anti-Hsp70 antibody. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1601-11, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig7:** Immunohistochemical analysis of paraffin-embedded human kidney tissue using anti-Hsp70 antibody. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1601-11, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




**Fig8:** Immunohistochemical analysis of paraffin-embedded human small intestine tissue using anti-Hsp70 antibody. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1601-11, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




**Fig9:** Immunohistochemical analysis of paraffin-embedded mouse testis tissue with Rabbit anti-Hsp70 antibody (ET1601-11) at 1/500 dilution.


The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1601-11) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




**Fig10:** Immunohistochemical analysis of paraffin-embedded mouse brain tissue using anti-Hsp70 antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1601-11, 1/100) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig11:** Immunohistochemical analysis of paraffin-embedded mouse prostate tissue using anti-Hsp70 antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1601-11, 1/100) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




**Fig12:** Flow cytometric analysis of Hsp70 was done on Hela cells. The cells were fixed, permeabilized and stained with the primary antibody (ET1601-11, 1/50) (blue). After incubation of the primary antibody at room temperature for an hour, the cells were stained with a Alexa Fluor 488-conjugated Goat anti-Rabbit IgG Secondary antibody at 1/1,000 dilution for 30 minutes. Unlabelled sample was used as a control (cells without incubation with primary antibody; red).



**Fig13:** Immunohistochemical analysis of paraffin-embedded rat prostate tissue with Rabbit anti-Hsp70 antibody (ET1601-11) at 1/1,000 dilution.


The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1601-11) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig14:** Immunohistochemical analysis of paraffin-embedded rat testis tissue with Rabbit anti-Hsp70 antibody (ET1601-11) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1601-11) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


**Fig15:** Immunocytochemistry analysis of C2C12 cells labeling Hsp70 with Rabbit anti-Hsp70 antibody (ET1601-11) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Hsp70 antibody (ET1601-11) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

**Fig16:** Immunocytochemistry analysis of NIH/3T3 cells labeling Hsp70 with Rabbit anti-Hsp70 antibody (ET1601-11) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Hsp70 antibody (ET1601-11) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

### Background References

1. "The disordered amino-terminus of SIMPL interacts with members of the 70-kDa heat-shock protein family." Haag Breese E., Uversky V.N., Georgiadis M.M., Harrington M.A. *DNA Cell Biol.* 25:704-714(2006)
2. "Hsp70 interacts with the retroviral restriction factor TRIM5alpha and assists the folding of TRIM5alpha." Hwang C.Y., Holl J., Rajan D., Lee Y., Kim S., Um M., Kwon K.S., Song B. *J. Biol. Chem.* 285:7827-7837(2010)

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn


 华安生物  
 HUABIO  
 www.huabio.cn