

# Anti-Smad2 Antibody [SP06-05]

ET1604-22



<b>Product Type:</b>	Recombinant Rabbit monoclonal IgG, primary antibodies
<b>Species reactivity:</b>	Human, Mouse, Rat
<b>Applications:</b>	WB, IF-Cell, IHC-P, IP, FC
<b>Molecular Wt:</b>	Predicted band size: 52 kDa
<b>Clone number:</b>	SP06-05

**Description:** Smad proteins, the mammalian homologs of the Drosophila mothers against decapentaplegic (Mad), have been implicated as downstream effectors of TGF $\beta$ /BMP signaling. Smad1 (also designated Madr1 or JV4-1) and Smad5 are effectors of BMP-2 and BMP-4 function, while Smad2 (also designated Madr2 or JV18-1) and Smad3 are involved in TGF $\beta$  and Activin-mediated growth modulation. Smad4 (also designated DPC4) has been shown to mediate all of the above activities through interaction with various Smad family members. Smad6 and Smad7 regulate the response to Activin/TGF $\beta$  signaling by interfering with TGF $\beta$ -mediated phosphorylation of other Smad proteins.

**Immunogen:** Synthetic peptide within human Smad2 aa 220-270.

**Positive control:** HeLa cell lysate, HT-29 cell lysate, Jurkat cell lysate, HL-60 cell lysate, C2C12 cell lysate, mouse lung tissue lysate, mouse placenta tissue lysate, human cerebellum tissue, mouse cerebellum tissue, rat cerebellum tissue, HepG2, NIH/3T3.

**Subcellular location:** Cytoplasm, Nucleus.

**Database links:** SwissProt: Q15796 Human | Q62432 Mouse | O70436 Rat

**Recommended Dilutions:**

<b>WB</b>	1:1,000-1:5,000
<b>IF-Cell</b>	1:100
<b>IHC-P</b>	1:200-1:1,000
<b>FC</b>	1:1,000
<b>IP</b>	Use at an assay dependent concentration.

**Storage Buffer:** 1\*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

**Storage Instruction:** Store at +4°C after thawing. Aliquot store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

**Purity:** Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn

## Images

**Fig1:** Western blot analysis of Smad2 on different lysates with Rabbit anti-Smad2 antibody (ET1604-22) at 1/5,000 dilution.

Lane 1: HeLa cell lysate  
Lane 2: HT-29 cell lysate  
Lane 3: Jurkat cell lysate  
Lane 4: HL-60 cell lysate



Lysates/proteins at 15 µg/Lane.

Predicted band size: 52 kDa  
Observed band size: 58 kDa


Exposure time: 1 minute 20 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDm/TBST for 1 hour at room temperature. The primary antibody (ET1604-22) at 1/5,000 dilution was used in 5% NFDm/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1:50,000 dilution was used for 1 hour at room temperature.

**Fig2:** Western blot analysis of Smad2 on different lysates with Rabbit anti-Smad2 antibody (ET1604-22) at 1/1,000 dilution.

Lane 1: Jurkat cell lysate (15 µg/Lane)  
Lane 2: C2C12 cell lysate (15 µg/Lane)  
Lane 3: Mouse lung tissue lysate (30 µg/Lane)  
Lane 4: Mouse placenta tissue lysate (30 µg/Lane)



Predicted band size: 52 kDa  
Observed band size: 58 kDa

Exposure time: 1 minute 21 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDm/TBST for 1 hour at room temperature. The primary antibody (ET1604-22) at 1/1,000 dilution was used in 5% NFDm/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.


Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn




**Fig3:** Western blot analysis of Smad2 with anti-Smad2 antibody (ET1604-22) at 1:500 dilution.

Lane 1: Wild-type HaCaT whole cell lysate (15  $\mu$ g).


Lane 2: Smad2 knockout HaCaT whole cell lysate (15  $\mu$ g).

ET1604-22 was shown to specifically react with Smad2 in wild-type HaCaT cells. NO band was observed when Smad2 knockout sample was tested. Wild-type and Smad2 knockout samples were subjected to SDS-PAGE. Proteins were transferred to a PVDF membrane and blocked with 5% NFDM in TBST for 1 hour at room temperature. The primary antibody (ET1604-22, 1:500) was used in 5% BSA at room temperature for 2 hours. Goat anti-Rabbit IgG-HRP antibody at 1:10,000 dilution was used for 1 hour at room temperature.




**Fig4:** Immunohistochemical analysis of paraffin-embedded human cerebellum tissue with Rabbit anti-Smad2 antibody (ET1604-22) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1604-22) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig5:** Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue with Rabbit anti-Smad2 antibody (ET1604-22) at 1/1,000 dilution.


The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1604-22) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig6:** Immunohistochemical analysis of paraffin-embedded rat cerebellum tissue with Rabbit anti-Smad2 antibody (ET1604-22) at 1/200 dilution.


The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1604-22) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

**Fig7:** Immunocytochemistry analysis of HepG2 cells labeling Smad2 with Rabbit anti-Smad2 antibody (ET1604-22) at 1/100 dilution.




Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Smad2 antibody (ET1604-22) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI. Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

**Fig8:** Immunocytochemistry analysis of NIH/3T3 cells labeling Smad2 with Rabbit anti-Smad2 antibody (ET1604-22) at 1/100 dilution.




Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Smad2 antibody (ET1604-22) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI. Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.



**Fig9:** Flow cytometric analysis of HepG2 cells labeling Smad2.

Cells were fixed and permeabilized. Then stained with the primary antibody (ET1604-22, 1/1,000) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4°C for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1121) at 1/1,000 dilution for 30 minutes at +4°C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).



**Fig10:** Flow cytometric analysis of NIH/3T3 cells labeling Smad2.

Cells were fixed and permeabilized. Then stained with the primary antibody (ET1604-22, 1/1,000) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4°C for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1121) at 1/1,000 dilution for 30 minutes at +4°C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).

**Note:** All products are “FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE”.

### Background References

1. Ungefroren H et al. Rac1b negatively regulates TGF-1-induced cell motility in pancreatic ductal epithelial cells by suppressing Smad signalling. *Oncotarget* 5:277-90 (2014).
2. Harazono Y et al. miR-655 Is an EMT-suppressive MicroRNA targeting ZEB1 and TGFBR2. *PLoS One* 8:e62757 (2013).

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
 HUABIO  
[www.huabio.cn](http://www.huabio.cn)