

Anti-BDNF Antibody [SJ12-09]

ET1606-42



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Mouse, Rat, Zebrafish
Applications:	WB, IF-Cell, FC
Molecular Wt:	Predicted band size: 28 kDa
Clone number:	SJ12-09

Description: BDNF acts on certain neurons of the central nervous system and the peripheral nervous system, helping to support survival of existing neurons, and encouraging growth and differentiation of new neurons and synapses. In the brain it is active in the hippocampus, cortex, and basal forebrain—areas vital to learning, memory, and higher thinking. BDNF is also expressed in the retina, kidneys, prostate, motor neurons, and skeletal muscle, and is also found in saliva. BDNF itself is important for long-term memory. Although the vast majority of neurons in the mammalian brain are formed prenatally, parts of the adult brain retain the ability to grow new neurons from neural stem cells in a process known as neurogenesis. Neurotrophins are proteins that help to stimulate and control neurogenesis, BDNF being one of the most active. Mice born without the ability to make BDNF have developmental defects in the brain and sensory nervous system, and usually die soon after birth, suggesting that BDNF plays an important role in normal neural development. Other important neurotrophins structurally related to BDNF include NT-3, NT-4, and NGF. BDNF is made in the endoplasmic reticulum and secreted from dense-core vesicles. It binds carboxypeptidase E (CPE), and disruption of this binding has been proposed to cause the loss of sorting BDNF into dense-core vesicles. The phenotype for BDNF knockout mice can be severe, including postnatal lethality. Other traits include sensory neuron losses that affect coordination, balance, hearing, taste, and breathing. Knockout mice also exhibit cerebellar abnormalities and an increase in the number of sympathetic neurons. Certain types of physical exercise have been shown to markedly (threefold) increase BDNF synthesis in the human brain, a phenomenon which is partly responsible for exercise-induced neurogenesis and improvements in cognitive function. Niacin appears to upregulate BDNF and tropomyosin receptor kinase B (TrkB) expression as well.

Immunogen:	Synthetic peptide within C-terminal human BDNF.
Positive control:	SH-SY5Y cell lysate, Neuro-2a cell lysate, mouse brain tissue lysate, rat brain tissue lysate, rat hippocampus tissue lysate, mouse hippocampus tissue lysate, Hela.
Subcellular location:	Secreted.
Database links:	SwissProt: P23560 Human P21237 Mouse P23363 Rat
Recommended Dilutions:	
WB	1:2,000-1:20,000
IF-Cell	1:50-1:200
FC	1:1,000
Storage Buffer:	1*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.
Storage Instruction:	Store at +4°C after thawing. Aliquot store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
Purity:	Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images

Fig1: Western blot analysis of BDNF on different lysates with Rabbit anti-BDNF antibody (ET1606-42) at 1/2,000 dilution.

Lane 1: SH-SY5Y cell lysate
 Lane 2: Neuro-2a cell lysate
 Lane 3: Mouse brain tissue lysate
 Lane 4: Rat brain tissue lysate
 Lane 5: Rat hippocampus tissue lysate
 Lane 6: Mouse hippocampus tissue lysate



Lysates/proteins at 10 µg/Lane.

Predicted band size: 28 kDa

Observed band size: 28 kDa


Exposure time: 1 minute 40 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody (ET1606-42) at 1/2,000 dilution was used in 5% NFDN/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig2: Western blot analysis of BDNF on different lysates with Rabbit anti-BDNF antibody (ET1606-42) at 1/20,000 dilution.

Lane 1: MDA-MB-231-si NT cell lysate
 Lane 2: MDA-MB-231-si BDNF cell lysate



Lysates/proteins at 10 µg/Lane.

Predicted band size: 28 kDa

Observed band size: 28 kDa

Exposure time: 14 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody (ET1606-42) at 1/20,000 dilution was used in 5% NFDN/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/100,000 dilution was used for 1 hour at room temperature.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
 HUABIO
 www.huabio.cn





Fig3: ICC staining of BDNF in HeLa cells (green). Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 1% Blocker BSA for 15 minutes at room temperature. Cells were probed with the primary antibody (ET1606-42, 1/200) for 1 hour at room temperature, washed with PBS. Alexa Fluor®488 Goat anti-Rabbit IgG was used as the secondary antibody at 1/1,000 dilution. The nuclear counter stain is DAPI (blue).

Fig4: Immunocytochemistry analysis of Neuro-2a cells labeling BDNF with Rabbit anti-BDNF antibody (ET1606-42) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-BDNF antibody (ET1606-42) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.




Fig5: Flow cytometric analysis of Neuro-2a cells labeling BDNF.

Cells were fixed and permeabilized. Then stained with the primary antibody (ET1606-42, 1µg/mL) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4°C for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1121) at 1/1,000 dilution for 30 minutes at +4°C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Adzic, M. et al. 2015. The contribution of hypothalamic neuroendocrine, neuroplastic and neuroinflammatory processes to lipopolysaccharide-induced depressive-like behaviour in female and male rats: Involvement of glucocorticoid receptor and C/EBP- β . Behavioural brain research. 291: 130-139.
2. Xu, Q. et al. 2015. Sigma 1 receptor activation regulates brain-derived neurotrophic factor through NR2A-CaMKIV-TORC1 pathway to rescue the impairment of learning and memory induced by brain ischaemia/reperfusion. Psychopharmacology. 232: 1779-91.

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
H U A B I O
www.huabio.cn