

Anti-LC3B Antibody [JJ090-6]

ET1701-65



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IF-Cell, IHC-P, IF-Tissue, IP
Molecular Wt:	Predicted band size: 14/16 kDa
Clone number:	JJ090-6

Description: Microtubule-associated proteins (MAPs) regulate microtubule stability and play critical roles in neuronal development and in maintaining the balance between neuronal plasticity and rigidity. MAP-light chain 3 beta (MAP-LC3 β) and MAP-light chain 3 alpha (MAP-LC3 α) are subunits of both MAP1A and MAP1B. MAP-LC3 β , a homolog of Apg8p, is essential for autophagy and associated to the autophagosome membranes after processing. Two forms of LC3 β , the cytosolic LC3-I and the membrane-bound LC3-II, are produced post-translationally. LC3-I is formed by the removal of the C-terminal 22 amino acids from newly synthesized LC3 β , followed by the conversion of a fraction of LC3-I into LC3-II. LC3 enhances fibronectin mRNA translation in ductus arteriosus cells through association with 60S ribosomes and binding to an AU-rich element in the 3' untranslated region of fibronectin mRNA. This facilitates sorting of fibronectin mRNA onto rough endoplasmic reticulum and translation. MAP LC3 β may also be involved in formation of autophagosomal vacuoles. It is expressed primarily in heart, testis, brain and skeletal muscle.

Immunogen: Synthetic peptide within human LC3 B aa 1-20 and aa 80-120.

Positive control: HeLa cells treated with 50 μ M Chloroquine for 24 hours, HeLa cell lysate, HeLa treated with 50 μ M Chloroquine for 18 hours cell lysate, C2C12 cell lysate, C2C12 treated with 50 μ M Chloroquine for 18 hours cell lysate, C6 cell lysate, C6 treated with 50 μ M Chloroquine for 18 hours cell lysate, mouse brain tissue lysate, rat brain tissue lysate, HCT 116 cell lysate, HCT 116 treated with 50 μ M Chloroquine for 18 hours cell lysate, U-87 MG cell lysate, C2C12 cells treated with 50 μ M Chloroquine for 24 hours, C6 cells treated with 50 μ M Chloroquine for 24 hours, mouse brain tissue, mouse hippocampus tissue, rat brain tissue, rat hippocampus tissue.

Subcellular location: Cytoplasm, Cytoplasmic vesicle, Cytoskeleton, Membrane, Microtubule, Mitochondrion.

Database links: SwissProt: Q9GZQ8 Human | Q9CQV6 Mouse | Q62625 Rat

Recommended Dilutions:

WB	1:1,000-1:2,000
IF-Cell	1:100-1:200
IHC-P	1:1,000-1:5,000
IF-Tissue	1:200-1:1,000
IP	Use at an assay dependent concentration.

Storage Buffer: 1*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Store at +4 $^{\circ}$ C after thawing. Aliquot store at -20 $^{\circ}$ C or -80 $^{\circ}$ C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images


Fig1: Immunocytochemistry analysis of HeLa cells treated with or without 50 μ M Chloroquine for 24 hours labeling LC3B with Rabbit anti-LC3B antibody (ET1701-65) at 1/200 dilution.



Cells were fixed in 4% paraformaldehyde for 10 minutes at 37 °C, permeabilized with 0.05% Triton X-100 in PBS for 20 minutes, and then blocked with 2% negative goat serum for 30 minutes at room temperature. Cells were then incubated with Rabbit anti-LC3B antibody (ET1701-65) at 1/200 dilution in 2% negative goat serum overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

Fig2: Western blot analysis of LC3B on different lysates with Rabbit anti-LC3B antibody (ET1701-65) at 1/2,000 dilution and competitor's antibody at 1/1,000 dilution.



Lane 1: HeLa cell lysate
 Lane 2: HeLa treated with 50 μ M Chloroquine for 18 hours cell lysate
 Lane 3: C2C12 cell lysate
 Lane 4: C2C12 treated with 50 μ M Chloroquine for 18 hours cell lysate
 Lane 5: C6 cell lysate
 Lane 6: C6 treated with 50 μ M Chloroquine for 18 hours cell lysate
 Lane 7: mouse brain tissue lysate
 Lane 8: rat brain tissue lysate
 Lysates/proteins at 20 μ g/Lane.

Predicted band size: 14/16 kDa
 Observed band size: 14/16 kDa

Exposure time: 3 minutes;
 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (ET1701-65) at 1/2,000 dilution and competitor's antibody at 1/1,000 dilution were used in 5% NFDM/TBST at 4 °C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
 HUABIO
 www.huabio.cn

Fig3: Western blot analysis of LC3B on different lysates with Rabbit anti-LC3B antibody (ET1701-65) at 1/1,000 dilution.

Lane 1: HCT 116 cell lysate
 Lane 2: HCT 116 treated with 50 μ M Chloroquine for 18 hours cell lysate
 Lane 3: U-87 MG cell lysate
 Lane 4: C6 cell lysate
 Lane 5: Mouse brain tissue lysate
 Lane 6: Rat brain tissue lysate



Lysates/proteins at 20 μ g/Lane.

Predicted band size: 14/16 kDa
 Observed band size: 14/16 kDa


Exposure time: 26 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (ET1701-65) at 1/1,000 dilution was used in 5% NFDM/TBST at 4 $^{\circ}$ C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig4: Western blot analysis of LC3B on different lysates with Rabbit anti-LC3B antibody (ET1701-65) at 1/2,000 dilution.

Lane 1: HeLa-si NT cell lysate
 Lane 2: HeLa-si LC3B cell lysate



Lysates/proteins at 10 μ g/Lane.

Predicted band size: 14/16 kDa
 Observed band size: 14/16 kDa

Exposure time: 1 minute;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (ET1701-65) at 1/2,000 dilution was used in 5% NFDM/TBST at 4 $^{\circ}$ C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
 HUABIO
 www.huabio.cn





Fig5: Immunocytochemistry analysis of C2C12 cells treated with or without 50 μ M Chloroquine for 24 hours labeling LC3B with Rabbit anti-LC3B antibody (ET1701-65) at 1/100 dilution.


Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-LC3B antibody (ET1701-65) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI. Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

Fig6: Immunocytochemistry analysis of C6 cells treated with or without 50 μ M Chloroquine for 24 hours labeling LC3B with Rabbit anti-LC3B antibody (ET1701-65) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-LC3B antibody (ET1701-65) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI. Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

Fig7: Immunohistochemical analysis of paraffin-embedded mouse brain tissue with Rabbit anti-LC3B antibody (ET1701-65) at 1/5,000 dilution.



The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ET1701-65) at 1/5,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn



Fig8: Immunohistochemical analysis of paraffin-embedded mouse hippocampus tissue with Rabbit anti-LC3B antibody (ET1701-65) at 1/5,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ET1701-65) at 1/5,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig9: Immunohistochemical analysis of paraffin-embedded rat brain tissue with Rabbit anti-LC3B antibody (ET1701-65) at 1/5,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ET1701-65) at 1/5,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig10: Immunohistochemical analysis of paraffin-embedded rat hippocampus tissue with Rabbit anti-LC3B antibody (ET1701-65) at 1/5,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ET1701-65) at 1/5,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Note: All products are “FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE”.

Background References

1. Gai Z et al. Farnesoid X Receptor Protects against Kidney Injury in Uninephrectomized Obese Mice. *J Biol Chem* 291:2397-411 (2016).
2. Xu TX et al. Hypoxia responsive miR-210 promotes cell survival and autophagy of endometriotic cells in hypoxia. *Eur Rev Med Pharmacol Sci* 20:399-406 (2016).

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
H U A B I O
www.huabio.cn