# Anti-Phospho-POLR2A (S2) Antibody [JM11-51]

# ET1703-86

Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IP, IF-Cell, IF-Tissue, IHC-P, FC
Molecular Wt:	Predicted band size: 217 kDa
Clone number:	JM11-51
Description:	DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Largest and catalytic component of RNA polymerase II which synthesizes mRNA precursors and many functional non-coding RNAs. Forms the polymerase active center together with the second largest subunit. Pol II is the central component of the basal RNA polymerase II transcription machinery. It is composed of mobile elements that move relative to each other. RPB1 is part of the core element with the central large cleft, the clamp element that moves to open and close the cleft and the jaws that are thought to grab the incoming DNA template. At the start of transcription, a single-stranded DNA template strand of the promoter is positioned within the central active site cleft of Pol II. A bridging helix emanates from RPB1 and crosses the cleft near the catalytic site and is thought to promote translocation of Pol II by acting as a ratchet that moves the RNA-DNA hybrid through the active site by switching from straight to bent conformations at each step of nucleotide addition. During transcription elongation, Pol II moves on the template as the transcript elongates. Elongation is influenced by the phosphorylation status of the C-terminal domain (CTD) of Pol II largest subunit (RPB1), which serves as a platform for assembly of factors that regulate transcription initiation, elongation, termination and mRNA processing. Acts as an RNA-dependent RNA polymerase when associated with small delta antigen of Hepatitis delta virus, acting both as a replicate and transcriptase for the viral RNA processing.
lmmunogen:	Synthetic phospho-peptide corresponding to residues surrounding Ser2 of Human POLR2A aa 1590-1630 / 1970.
Positive control:	HeLa cell lysate, RAW264.7 cell lysate, PC-12 cell lysate, HeLa, RAW264.7, PC-12, human testis tissue, mouse testis tissue, rat testis tissue, 商品名 was done on Hela.
Subcellular location:	Nucleus.
Database links:	SwissProt: P24928 Human   P08775 Mouse Entrez Gene: 363633 Rat
Recommended Dilutions: WB IF-Cell IF-Tissue IHC-P FC Storage Buffer: Storage Instruction:	1:500-1:2,000 1:50-1:100 1:100-1:500 1:1,000 1:1,000 1*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide. Shipped at $4^{\circ}$ C. Store at $+4^{\circ}$ C short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at $-20^{\circ}$ C long term.
Purity:	Protein A affinity purified.

# Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn



11.

#### Images



**Fig1:** Western blot analysis of Phospho-POLR2A (S2) on different lysates with Rabbit anti-Phospho-POLR2A (S2) antibody (ET1703-86) at 1/1,000 dilution.

Lane 1: HeLa cell lysate Lane 2: RAW264.7 cell lysate Lane 3: PC-12 cell lysate Lane 4: HeLa cell lysate, the membrane treated with  $\lambda pp$  for 1 hour

Lysates/proteins at 20 µg/Lane.

Predicted band size: 217 kDa Observed band size: 260 kDa

Exposure time: 6 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (ET1703-86) at 1/1,000 dilution was used in 5% NFDM/TBST at  $4^{\circ}$ C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

**Fig2:** Immunocytochemistry analysis of HeLa cells labeling Phospho-POLR2A (S2) with Rabbit anti-Phospho-POLR2A (S2) antibody (ET1703-86) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 15 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 15 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Phospho-POLR2A (S2) antibody (ET1703-86) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (HA601187, red) was stained at 1/100 dilution overnight at  $+4^{\circ}$ C. Goat Anti-Mouse IgG H&L (iFluor  $\pm$  594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

### Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn





**Fig3:** Immunocytochemistry analysis of RAW264.7 cells labeling Phospho-POLR2A (S2) with Rabbit anti-Phospho-POLR2A (S2) antibody (ET1703-86) at 1/50 dilution.

Cells were fixed in 4% paraformaldehyde for 15 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 15 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Phospho-POLR2A (S2) antibody (ET1703-86) at 1/50 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (HA601187, red) was stained at 1/100 dilution overnight at +4 $^{\circ}$ C. Goat Anti-Mouse IgG H&L (iFluor <sup>TM</sup> 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.



**Fig4:** Immunocytochemistry analysis of PC-12 cells labeling Phospho-POLR2A (S2) with Rabbit anti-Phospho-POLR2A (S2) antibody (ET1703-86) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 15 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 15 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Phospho-POLR2A (S2) antibody (ET1703-86) at 1/100 dilution in 1% BSA in PBST overnight at 4  $^{\circ}$ C. Goat Anti-Rabbit IgG H&L (iFluor TM 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (HA601187, red) was stained at 1/100 dilution overnight at +4℃. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

### Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn





**Fig5:** Immunohistochemical analysis of paraffin-embedded human testis tissue untreated / treated with  $\lambda$ pp with Rabbit anti-Phospho-POLR2A (S2) antibody (ET1703-86) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1703-86) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

**Fig6:** Immunohistochemical analysis of paraffin-embedded mouse testis tissue untreated / treated with  $\lambda pp$  with Rabbit anti-Phospho-POLR2A (S2) antibody (ET1703-86) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1703-86) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

**Fig7:** Immunohistochemical analysis of paraffin-embedded rat testis tissue untreated / treated with  $\lambda$ pp with Rabbit anti-Phospho-POLR2A (S2) antibody (ET1703-86) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1703-86) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

**Fig8:** Flow cytometric analysis of Phospho-POLR2A (S2) was done on Hela cells. The cells were fixed, permeabilized and stained with the primary antibody (ET1703-86, 1/50) (red). After incubation of the primary antibody at room temperature for an hour, the cells were stained with a Alexa Fluor®488 conjugate-Goat anti-Rabbit IgG Secondary antibody at 1/1,000 dilution for 30 minutes.Unlabelled sample was used as a control (cells without incubation with primary antibody; black).

### Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

**が作安生物** www.huabio.cn





Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

#### **Background References**

- 1. Rother S et al. NF-kB-repressing factor phosphorylation regulates transcription elongation via its interactions with 5'-3' exoribonuclease 2 and negative elongation factor. FASEB J 30:174-85 (2016).
- 2. Ard R & Allshire RC Transcription-coupled changes to chromatin underpin gene silencing by transcriptional interference. Nucleic Acids Res N/A:N/A (2016).

### Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880



Service mail:support@huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

Technical:0086-571-89986345