

Anti-hnRNP Q Antibody [JG35-72]

ET7108-17



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Rat, Mouse
Applications:	WB, IHC-P, IF-Cell, IF-Tissue
Molecular Wt:	Predicted band size: 70 kDa
Clone number:	JG35-72

Description: Pre-mRNA splicing is a critical step in the posttranscriptional regulation of gene expression. Heterogeneous nuclear ribonucleoprotein Q (hnRNP Q) is involved in RNA processing and is necessary for efficient pre-mRNA splicing. hnRNP is widely expressed and developmentally regulated. hnRNP Q interacts with survival motor neuron protein (SMN). Loss of function of SMN results in spinal muscular atrophy, a common neurodegenerative disease. The most common deletion in SMN genes disrupts the interaction between SMN and hnRNP Q. hnRNP Q is upregulated after midnight, and this upregulation correlates with an abrupt decline in AANAT, the key enzyme in melatonin synthesis. Rhythmic AANAT mRNA degradation mediated in part by hnRNP Q implicates this enzyme in the regulation of circadian oscillation.

Immunogen: Synthetic peptide within Human hnRNP Q aa 574-623 / 623.

Positive control: K562 cell lysates, rat brain tissue lysates, LOVO, SiHa, rat testis tissue, human thyroid tissue, human breast tissue, human kidney tissue.

Subcellular location: Cytoplasm, Endoplasmic reticulum, Microsome, Nucleus, Spliceosome.

Database links: SwissProt: O60506 Human | Q7TP47 Rat | Q7TMK9 Mouse

Recommended Dilutions:

WB	1:500-1:2,000
IF-Cell	1:50-1:200
IF-Tissue	1:50-1:200
IHC-P	1:50-1:200

Storage Buffer: 1*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images




Fig1: Western blot analysis of hnRNP Q on K562 cell lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (ET7108-17, 1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1:200,000 dilution was used for 1 hour at room temperature.




Fig2: Western blot analysis of hnRNP Q on rat brain tissue lysates with Rabbit anti-hnRNP Q antibody (ET7108-17) at 1/500 dilution.

Lysates/proteins at 20 µg/Lane.

Predicted band size: 70 kDa

Observed band size: 70 kDa

Exposure time: 2 minutes;

8% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody (ET7108-17) at 1/500 dilution was used in 5% NFDN/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1:200,000 dilution was used for 1 hour at room temperature.




Fig3: ICC staining of hnRNP Q in LOVO cells (green). Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 10% negative goat serum for 15 minutes at room temperature. Cells were probed with the primary antibody (ET7108-17, 1/50) for 1 hour at room temperature, washed with PBS. Alexa Fluor®488 conjugate-Goat anti-Rabbit IgG was used as the secondary antibody at 1/1,000 dilution. The nuclear counter stain is DAPI (blue).




Fig4: ICC staining of hnRNP Q in SiHa cells (green). Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 10% negative goat serum for 15 minutes at room temperature. Cells were probed with the primary antibody (ET7108-17, 1/50) for 1 hour at room temperature, washed with PBS. Alexa Fluor®488 conjugate-Goat anti-Rabbit IgG was used as the secondary antibody at 1/1,000 dilution. The nuclear counter stain is DAPI (blue).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn




Fig5: Immunohistochemical analysis of paraffin-embedded rat testis tissue using anti-hnRNP Q antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ET7108-17, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig6: Immunohistochemical analysis of paraffin-embedded human thyroid tissue using anti-hnRNP Q antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ET7108-17, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig7: Immunohistochemical analysis of paraffin-embedded human kidney tissue using anti-hnRNP Q antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ET7108-17, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig8: Immunohistochemical analysis of paraffin-embedded human breast tissue with Rabbit anti-hnRNP Q antibody (ET7108-17) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ET7108-17) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Mourelatos Z et al. SMN interacts with a novel family of hnRNP and spliceosomal proteins. *EMBO J* 20:5443-5452 (2001).
2. Grosset C et al. A mechanism for translationally coupled mRNA turnover: interaction between the poly(A) tail and a c-fos RNA coding determinant via a protein complex. *Cell* 103:29-40 (2000).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUAABIO
www.huabio.cn