

# Anti-CD4 Antibody [JE56-36]

ET7110-64



<b>Product Type:</b>	Recombinant Rabbit monoclonal IgG, primary antibodies
<b>Species reactivity:</b>	Human
<b>Applications:</b>	WB, IHC-P
<b>Molecular Wt:</b>	51 kDa
<b>Clone number:</b>	JE56-36

**Description:** Integral membrane glycoprotein that plays an essential role in the immune response and serves multiple functions in responses against both external and internal offenses. In T-cells, functions primarily as a coreceptor for MHC class II molecule:peptide complex. The antigens presented by class II peptides are derived from extracellular proteins while class I peptides are derived from cytosolic proteins. Interacts simultaneously with the T-cell receptor (TCR) and the MHC class II presented by antigen presenting cells (APCs). In turn, recruits the Src kinase LCK to the vicinity of the TCR-CD3 complex. LCK then initiates different intracellular signaling pathways by phosphorylating various substrates ultimately leading to lymphokine production, motility, adhesion and activation of T-helper cells. In other cells such as macrophages or NK cells, plays a role in differentiation/activation, cytokine expression and cell migration in a TCR/LCK-independent pathway. Participates in the development of T-helper cells in the thymus and triggers the differentiation of monocytes into functional mature macrophages. (Microbial infection) Primary receptor for human immunodeficiency virus-1 (HIV-1). Acts as a receptor for Human Herpes virus 7/HHV-7.

**Immunogen:** Synthetic peptide within Human CD4 aa 409-458 / 458.

**Positive control:** THP-1 cell lysate, HL-60 cell lysate, human tonsil tissue, human spleen tissue.

**Subcellular location:** Cell membrane.

**Database links:** SwissProt: P01730 Human

**Recommended Dilutions:**

<b>WB</b>	1:500-1:2,000
<b>IHC-P</b>	1:50-1:200

**Storage Buffer:** 1\*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

**Storage Instruction:** Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

**Purity:** Protein A affinity purified.

Hangzhou HuaAn Biotechnology Co.,Ltd.

Orders: 0086-571-88062880

Technical:0086-571-89986345

Service mail: support@huabio.cn

www.huabio.cn



## Images

**Fig1:** Western blot analysis of CD4 on different lysates with Rabbit anti-CD4 antibody (ET7110-64) at 1/500 dilution.

Lane 1: THP-1 cell lysate


Lane 2: HL-60 cell lysate

Lysates/proteins at 10 µg/Lane.

Predicted band size: 51 kDa


Observed band size: 55 kDa

Exposure time: 2 minutes;




Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody (ET7110-64) at 1/500 dilution was used in 5% NFDN/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1:5,000 dilution was used for 1 hour at room temperature.

**Fig2:** Immunohistochemical analysis of paraffin-embedded human tonsil tissue with Rabbit anti-CD4 antibody (ET7110-64) at 1/200 dilution.



The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET7110-64) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

**Fig3:** Immunohistochemical analysis of paraffin-embedded human spleen tissue with Rabbit anti-CD4 antibody (ET7110-64) at 1/50 dilution.



The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET7110-64) at 1/50 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

---

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

---

### Background References

1. Sanchez-Martinez A. et. al. Cytotoxic CD4+ T-cells during HIV infection: Targets or weapons? J Clin Virol. 2019 Oct;119:17-23.
2. Tompa A. et. al. Subsets of CD4+, CD8+, and CD25hi Lymphocytes Are in General Not Influenced by Isolation and Long-Term Cryopreservation. J Immunol. 2018 Sep 15;201(6):1799-1809.

Hangzhou HuaAn Biotechnology Co.,Ltd.

Orders: 0086-571-88062880

Technical:0086-571-89986345

Service mail: support@huabio.cn

www.huabio.cn

