Anti-EGR1 Antibody [JE57-62]

ET7111-41



Product Type: Recombinant Rabbit monoclonal IgG, primary antibodies

Species reactivity: Human

Applications: WB, IHC-P

Molecular Wt: Predicted band size: 58 kDa.

Clone number: JE57-62

Description: The protein encoded by this gene belongs to the EGR family of C2H2-type zinc-finger

proteins. It is a nuclear protein and functions as a transcriptional regulator. The products of target genes it activates are required for differentitation and mitogenesis. Studies suggest

this is a cancer suppressor gene.

Immunogen: Recombinant protein within human EGR1 aa 1-200.

Positive control: THP-1 treated with 100ng/mL LPS for 3 hours cell lysate, human brain tissue, human thyroid

carcinoma tissue.

Subcellular location: Nucleus, Cytoplasm.

Database links: SwissProt: P18146 Human

Recommended Dilutions:

WB 1:1,000 **IHC-P** 1:200

Storage Buffer: 1*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Store at $+4^{\circ}$ C after thawing. Aliquot store at -20° C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.



Service mail:support@huabio.cn www.huabio.cn

Images

Fig1: Western blot analysis of EGR1 on different lysates with Rabbit anti-EGR1 antibody (ET7111-41) at 1/1,000 dilution.

Lane 1: THP-1 cell lysate

Lane 2: THP-1 treated with 100ng/mL LPS for 3 hours cell lysate

Lysates/proteins at 20 µg/Lane.

Predicted band size: 58 kDa Observed band size: 75 kDa

Exposure time: 1 minute; ECL: K1801;

4-20% SDS-PAGE gel.



Fig2: Immunohistochemical analysis of paraffin-embedded human brain tissue with Rabbit anti-EGR1 antibody (ET7111-41) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH $_2$ O and PBS, and then probed with the primary antibody (ET7111-41) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig3: Immunohistochemical analysis of paraffin-embedded human thyroid carcinoma tissue with Rabbit anti-EGR1 antibody (ET7111-41) at 1/800 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH $_2$ O and PBS, and then probed with the primary antibody (ET7111-41) at 1/800 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Hangzhou Huaan Biotechnology Co., Ltd.

// 华安生物 www.huabio.cn Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

- 1. Li L. et. al. EGR1 regulates angiogenic and osteoclastogenic factors in prostate cancer and promotes metastasis. Oncogene. 2019 Aug
- 2. Mohtar O. et. al. Egr1 mediates the effect of insulin on leptin transcription in adipocytes. J Biol Chem. 2019 Apr