

# Human Tim-3/HAVCR2 Protein

HA210253



<b>Product name:</b>	Human Tim-3/HAVCR2 Protein
<b>Species reactivity:</b>	Human
<b>Protein construction description:</b>	Recombinant Human Tim-3/HAVCR2 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Ser22-Arg200.
<b>Background:</b>	Hepatitis A virus cellular receptor 2 (HAVCR2), also known as T-cell immunoglobulin and mucin-domain containing-3 (TIM-3), is a protein that in humans is encoded by the HAVCR2 gene. TIM3 is an immune checkpoint and together with other inhibitory receptors including programmed cell death protein 1 (PD-1) and lymphocyte activation gene 3 protein (LAG3) mediate the CD8 T-cell exhaustion. TIM3 has also been shown as a CD4 Th1-specific cell surface protein that regulates macrophage activation and enhances the severity of experimental autoimmune encephalomyelitis in mice.
<b>Purity:</b>	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC
<b>Endotoxin:</b>	Less than 1EU per µg by the LAL method.
<b>Source:</b>	HEK293
<b>Accession:</b>	Q8TDQ0-1
<b>Predicted molecular mass:</b>	The protein has a predicted MW of 20.7 kDa, Due to glycosylation, the protein migrates to 50-60 kDa based on Tris-Bis PAGE result
<b>Formulation:</b>	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
<b>Reconstitution:</b>	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
<b>Storage:</b>	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn


 华安生物  
HUABIO  
www.huabio.cn



**Fig1:** Human TIM-3 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



**Fig2:** The purity of Human TIM-3 is greater than 95% as determined by SEC-HPLC.



**Fig3:** Immobilized Human TIM-3, His Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Anti-TIM-3 Antibody, hFc Tag with the EC50 of 4.5ng/ml determined by ELISA.

**Note:** All products are “FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE”.