Human Mesothelin, C-His Tag Protein HA210521



Product name:	Human Mesothelin, C-His Tag			
Species reactivity:	Human			
Bio-Activity:	Testing in progress.			
Protein construction description:	A DNA sequence encoding the human Mesothelin protein (Q13421-3) (Glu 296-Ser 598) was expressed with a His at the C-terminus.			
Background:	Mesothelin, also known as MSLN, is a protein that in humans is encoded by the MSLN gene. Mesothelin is a 40 kDa protein that is expressed in mesothelial cells. The protein was first identified by its reactivity with monoclonal antibody K1. Subsequent cloning studies showed that the mesothelin gene encodes a precursor protein that is processed to yield mesothelin which is attached to the cell membrane by a glycophosphatidylinositol linkage and a 31-kDa shed fragment named megakaryocyte-potentiating factor (MPF). Although it has been proposed that mesothelin may be involved in cell adhesion, its biological function is not known. A knockout mouse line that lacks mesothelin reproduces and develops normally. Mesothelin is over expressed in several human tumors, including mesothelioma, ovarian cancer, pancreatic adenocarcinoma, lung adenocarcinoma, and cholangiocarcinoma. Mesothelin binds MUC16 (also known as CA125), indicating that the interaction of mesothelin and MUC16 may contribute to the implantation and peritoneal spread of tumors by cell adhesion. The region (residues 296-359) consisting of 64 amino acids at the N-terminus of cell surface mesothelin has been identified as the functional binding domain (named IAB) for MUC16/CA125, suggesting the mechanism of mesothelin acting as a MUC16/CA125 functional partner in cancer development.			
Purity:	>95% as determined by SDS-PAGE.			
Endotoxin:	Less than 1.0 EU per μ g by the LAL method.			
Fragment region:	Mesothelin (296-598)			
Source:	HEK293			
Accession:	Q13421-3			
Predicted molecular mass:	35.5 kD			
Formulation:	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4, 5% Trehalose, 5% mannitol.			
Reconstitution:	stitution: Reconstitute at 250 µg/ml in sterile water.			
Storage:	Please avoid repeated freeze-thaw cycles. Samples are stable for up to twelve months from date of receipt at - 20° C to - 80° C It is recommended that aliquot the reconstituted solution to minimize freeze-thaw cycles.			

Hangzhou Huaan Biotechnology Co., Ltd.

Ord	larc'	006_	571_	00061	000
JIU	lers.u	000-	571-	0000	2000

Technical:0086-571-89986345

Service mail:support@huabio.cn



Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

Images



Fig1: Protein on SDS-PAGE under reducing (R) condition.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Hangzhou Huaan Biotechnology Co., Ltd.



Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation