Human CD2, C-His, Flag Tag (ECD) Protein HA210554



Product name: Human CD2, C-His, Flag Tag (ECD)

Species reactivity: Human

Bio-Activity: Testing in progress.

Protein construction

description:

A DNA sequence encoding the human CD2 protein (P06729) (Lys 25-Asp 209) was expressed with both His

and Flag tag at the C-terminus.

Background: The protein encoded by this gene is a surface antigen found on all peripheral blood T-cells. The encoded

protein interacts with LFA3 (CD58) on antigen presenting cells to optimize immune recognition. A locus control region (LCR) has been found in the 3' flanking sequence of this gene. CD2 interacts with lymphocyte function-associated antigen CD58 (LFA-3) and CD48/BCM1 to mediate adhesion between T-cells and other cell types. CD2 is implicated in the triggering of T-cells, the cytoplasmic domain is implicated in the signaling function.

Purity: >95% as determined by SDS-PAGE.

Endotoxin: Less than 1.0 EU per µg by the LAL method.

Fragment region: CD2 (25-209)

Source: HEK293

Accession: P06729

Predicted molecular mass: 23.9 kD

Formulation: Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4, 5% Trehalose, 5% mannitol.

Reconstitution: Reconstitute at 250 µg/ml in sterile water.

Storage: Please avoid repeated freeze-thaw cycles. Samples are stable for up to twelve months from date of receipt at -

 20° to -80° It is recommended that aliquot the reconstituted solution to minimize freeze-thaw cycles.

Hangzhou Huaan Biotechnology Co., Ltd.

Technical: 0086-571-89986345

Service mail:support@huabio.cn



Images




Fig1: Protein on SDS-PAGE under reducing (R) condition.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".