

Mouse IL-1 beta/IL1B, C-His, Flag Tag Protein

HA210706



Product name:	Mouse IL-1 beta/IL1B, C-His, Flag Tag
Species reactivity:	Mouse
Bio-Activity:	Testing in progress.
Protein construction description:	A DNA sequence encoding the mouse IL-1 beta/IL1B protein (P10749) (Val 118-Ser 269) was expressed with both His, Flag tag at the C-terminus.

Background:	Potent pro-inflammatory cytokine. Initially discovered as the major endogenous pyrogen, induces prostaglandin synthesis, neutrophil influx and activation, T-cell activation and cytokine production, B-cell activation and antibody production, and fibroblast proliferation and collagen production. Promotes Th17 differentiation of T-cells. Synergizes with IL12/interleukin-12 to induce IFNG synthesis from T-helper 1 (Th1) cells. Plays a role in angiogenesis by inducing VEGF production synergistically with TNF and IL6. Involved in transduction of inflammation downstream of pyroptosis: its mature form is specifically released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore. IL1B production occurs in 2 steps, each being controlled by different stimuli. First, inflammatory signals, such as LPS, stimulate the synthesis and promote the accumulation of cytosolic stores of pro-IL1B (priming). Then additional signals are required for inflammasome assembly, leading to CASP1 activation, pro-IL1B processing and eventually secretion of the active cytokine. IL1B processing and secretion are temporarily associated.
--------------------	---

Purity:	>95% as determined by SDS-PAGE.
Endotoxin:	Less than 1.0 EU per µg by the LAL method.
Fragment region:	IL-1 beta/IL1B (118-269)
Source:	E.coli
Accession:	P10749
Predicted molecular mass:	20.7 kD
Formulation:	Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4, 5% Trehalose, 5% mannitol.
Reconstitution:	Reconstitute at 250 µg/ml in sterile water.
Storage:	Please avoid repeated freeze-thaw cycles. Samples are stable for up to twelve months from date of receipt at -20°C to -80°C. It is recommended that aliquot the reconstituted solution to minimize freeze-thaw cycles.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders: 0086-571-88062880

Technical: 0086-571-89986345

Service mail: support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Applications: WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation




Fig1: Protein on SDS-PAGE under reducing (R) condition.

Note: All products are “FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE”.