Human NCAM1/CD56, Tag Free (ECD) Protein HA210820



Product name:	Human NCAM1/CD56, Tag Free (ECD)									
Species reactivity:	Human									
Bio-Activity:	Testing in progress.									
Protein construction description:	A DNA sequence encoding the human NCAM1/CD56 protein (P13591-2) (Leu 20-Gly 718) was expressed with tag free.									
Background:	This gene encodes a cell adhesion protein which is a member of the immunoglobulin superfamily. The encoded protein is involved in cell-to-cell interactions as well as cell-matrix interactions during development and differentiation. The encoded protein plays a role in the development of the nervous system by regulating neurogenesis, neurite outgrowth, and cell migration. This protein is also involved in the expansion of T lymphocytes, B lymphocytes and natural killer (NK) cells which play an important role in immune surveillance. This protein plays a role in signal transduction by interacting with fibroblast growth factor receptors, N-cadherin and other components of the extracellular matrix and by triggering signalling cascades involving FYN-focal adhesion kinase (FAK), mitogen-activated protein kinase (MAPK), and phosphatidylinositol 3-kinase (PI3K). One prominent isoform of this gene, cell surface molecule CD56, plays a role in several myeloproliferative disorders such as acute myeloid leukemia and differential expression of this gene is associated with differential disease progression. For example, increased expression of CD56 is correlated with lower survival in acute myeloid leukemia patients whereas increased severity of COVID-19 is correlated with decreased abundance of CD56-expressing NK cells in peripheral blood. Alternative splicing results in multiple transcript variants encoding distinct protein isoforms.									
Purity:	>95% as determined by SDS-PAGE.									
Endotoxin:	Less than 1.0 EU per μ g by the LAL method.									
Fragment region:	NCAM1/CD56 (20-718)									
Source:	HEK293									
Accession:	P13591-2									
Predicted molecular mass:	77.7 kD									
Formulation:	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4, 5% Trehalose, 5% mannitol.									
Reconstitution:	Reconstitute at 250 µg/ml in sterile water.									
Storage:	Please avoid repeated freeze-thaw cycles. Samples are stable for up to twelve months from date of receipt at - 20° C to - 80° C It is recommended that aliquot the reconstituted solution to minimize freeze-thaw cycles.									

Hangzhou Huaan Biotechnology Co., Ltd.

0	rd	e	rs'I	00	8	6-	-57	71	-	88	0	62	88	30	
u	10			00	0	U	5			00	U	02	.00	50	

Technical:0086-571-89986345

5 Service mail:support@huabio.cn



Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

Images



Fig1: Protein on SDS-PAGE under reducing (R) condition.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Hangzhou Huaan Biotechnology Co., Ltd.



Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation