

Human TREM1, C-Hia, Flag Tag Protein

HA210846



| | |
|--|--|
| Product name: | Human TREM1, C-Hia, Flag Tag |
| Species reactivity: | Human |
| Bio-Activity: | Testing in progress. |
| Protein construction description: | A DNA sequence encoding the human TREM1 protein (Q9NP99-1) (Ala 21-Asn 205) was expressed with both His, Flag tag at the C-terminus. |

Background: This gene encodes a receptor belonging to the Ig superfamily that is expressed on myeloid cells. This protein amplifies neutrophil and monocyte-mediated inflammatory responses triggered by bacterial and fungal infections by stimulating release of pro-inflammatory chemokines and cytokines, as well as increased surface expression of cell activation markers. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene. Cell surface receptor that plays important roles in innate and adaptive immunity by amplifying inflammatory responses. Upon activation by various ligands such as PGLYRP1, HMGB1 or HSP70, multimerizes and forms a complex with transmembrane adapter TYROBP/DAP12. In turn, initiates a SYK-mediated cascade of tyrosine phosphorylation, activating multiple downstream mediators such as BTK, MAPK1, MAPK3 or phospholipase C-gamma. This cascade promotes the neutrophil- and macrophage-mediated release of pro-inflammatory cytokines and/or chemokines, as well as their migration and thereby amplifies inflammatory responses that are triggered by bacterial and fungal infections. By also promoting the amplification of inflammatory signals that are initially triggered by Toll-like receptor (TLR) and NOD-like receptor engagement, plays a major role in the pathophysiology of acute and chronic inflammatory diseases of different etiologies including septic shock and atherosclerosis.

| | |
|----------------------------------|--|
| Purity: | >95% as determined by SDS-PAGE. |
| Endotoxin: | Less than 1.0 EU per µg by the LAL method. |
| Fragment region: | TREM1 (21-205) |
| Source: | HEK293 |
| Accession: | Q9NP99-1 |
| Predicted molecular mass: | 23.5 kD |
| Formulation: | Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4, 5% Trehalose, 5% mannitol. |
| Reconstitution: | Reconstitute at 250 µg/ml in sterile water. |
| Storage: | Please avoid repeated freeze-thaw cycles. Samples are stable for up to twelve months from date of receipt at -20°C to -80°C. It is recommended that aliquot the reconstituted solution to minimize freeze-thaw cycles. |

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn




Fig1: Protein on SDS-PAGE under reducing (R) condition.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".