Human IL-16, C-His Tag Protein HA210917



Product name:	Human IL-16, C-His Tag			
Species reactivity:	uman			
Bio-Activity:	Testing in progress.			
Protein construction description:	NA sequence encoding the human IL-16 protein (Q14005-1) (Met 1203-Ser 1332) was expressed with a ag at the C-terminus			
Background:	The cytokine encoded by this gene is a pleiotropic cytokine that functions as a chemoattractant, a modulator of cell activation, and an inhibitor of HIV replication. The signaling process of this cytokine is mediated by CD4. The product of this gene undergoes proteolytic processing, which is found to yield two functional proteins. The cytokine function is exclusively attributed to the secreted C-terminal peptide, while the N-terminal product material a role in cell cycle control. Caspase 3 is reported to be involved in the proteolytic processing of this protein Two alternatively spliced transcript variants encoding distinct isoforms have been reported. Interleukin 16 (IL 16) is released by a variety of cells (including lymphocytes and some epithelial cells) that has been characterized as a chemoattractant for certain immune cells expressing the cell surface molecule CD4. IL-16 was originally described as a factor that could attract activated T cells in humans, it was previously called lymphocy chemoattractant factor (LCF). Since then, this interleukin has been shown to recruit and activate many other cell expressing the CD4 molecule, including monocytes, eosinophils, and dendritic cells. The structure of IL-16 was determined following its cloning in 1994. This cytokine is produced as a precursor peptide (pro-IL-16) th requires processing by an enzyme called caspase-3 to become active. CD4 is the cell signaling receptor for mature IL-16.			
Purity:	>95% as determined by SDS-PAGE.			
Endotoxin:	Less than 1.0 EU per μ g by the LAL method.			
Fragment region:	IL-16 (1203-1332)			
Source:	HEK293			
Accession:	Q14005-1			
Predicted molecular mass:	14.8 kD			
Formulation:	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4, 5% Trehalose, 5% mannitol.			
Reconstitution:	Reconstitute at 250 µg/ml in sterile water.			
Storage:	Please avoid repeated freeze-thaw cycles. Samples are stable for up to twelve months from date of receipt at - 20° C to - 80° C It is recommended that aliquot the reconstituted solution to minimize freeze-thaw cycles.			

Hangzhou Huaan Biotechnology Co., Ltd.

Ord	larc'	006_	571_	00061	000
JIU	lers.u	000-	571-	0000	2000

Technical:0086-571-89986345

5 Service mail:support@huabio.cn



Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

Images



Fig1: Protein on SDS-PAGE under reducing (R) condition.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Hangzhou Huaan Biotechnology Co., Ltd.



Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation