Anti-SFT Antibody HA500229



| Product Type: | Rabbit polyclonal IgG, primary antibodies |
|---------------------------------------|---|
| Species reactivity: | Human, Mouse, Rat |
| Applications: | WB, IHC-P |
| Molecular Wt: | 17 kDa |
| Description: | The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. This enzyme is closely related to a stimulator of iron transport (SFT), and is up-regulated in hereditary hemochromatosis. It also functions in the ubiquitination of the tumor-suppressor protein p53 and the hypoxia-inducible transcription factor HIF1alpha by interacting with the E1 ubiquitin-activating enzyme and the E3 ubiquitin-protein ligases. Two transcript variants encoding different isoforms have been found for this gene. |
| lmmunogen: | Recombinant protein within human SFT aa 1-147. |
| Positive control: | Hela cell lysate, Jurkat cell lysate, NIH/3T3 cell lysate, rat skeletal muscle tissue lysate, mouse liver tissue lysate, rat kidney tissue, human breast tissue, mouse skeletal muscle tissue. |
| Subcellular location: | Cytoplasm. |
| Database links: | SwissProt: P51668 Human P61080 Mouse D3ZDK2 Rat |
| Recommended Dilutions: WB IHC-P | 1:500-1:2,000 1:400 |
| Storage Buffer: | 1*TBS (pH7.4), 0.2% BSA, 50% Glycerol. Preservative: 0.05% Sodium Azide. |
| Storage Instruction: | Shipped at 4 $^\circ\!\!\mathbb{C}$. Store at +4 $^\circ\!\!\mathbb{C}$ short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20 $^\circ\!\!\mathbb{C}$ long term. |
| Purity: | Immunogen affinity purified. |

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

45 Service mail:support@huabio.cn



Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

HA500229 - Page 2

Images



Fig1: Western blot analysis of SFT on different lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (HA500229, 1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1:200,000 dilution was used for 1 hour at room temperature.

Positive control:

Lane 1: Hela cell lysate Lane 2: Jurkat cell lysate Lane 3: NIH/3T3 cell lysate Lane 4: Rat skeletal muscle tissue lysate Lane 5: Mouse liver tissue lysate





Fig2: Immunohistochemical analysis of paraffin-embedded rat kidney tissue using anti-SFT antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA

Fig3: Immunohistochemical analysis of paraffin-embedded human breast tissue using anti-SFT antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA500229, 1/400) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Fig4: Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue using anti-SFT antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA500229, 1/400) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn



Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

- 1. Zhou C. et. al. Gain of UBE2D1 facilitates hepatocellular carcinoma progression and is associated with DNA damage caused by continuous IL-6. J Exp Clin Cancer Res. 2018 Nov
- 2. Hou L. et. al. UBE2D1 RNA Expression Was an Independent Unfavorable Prognostic Indicator in Lung Adenocarcinoma, but Not in Lung Squamous Cell Carcinoma. Dis Markers. 2018 Oct

Hangzhou Huaan Biotechnology Co., Ltd.



Orders:0086-571-88062880

Technical:0086-571-89986345

5 Service mail:support@huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cetl=Immunofluorescence (Cetl) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation