

# Anti-Histone H3 (acetyl K27) Antibody [A6D6]

## HA600048



<b>Product Type:</b>	Mouse monoclonal IgG1, primary antibodies
<b>Species reactivity:</b>	Human, Mouse, Rat
<b>Applications:</b>	WB, IF-Cell, IHC-P, FC, ChIP
<b>Molecular Wt:</b>	Predicted band size: 15 kDa
<b>Clone number:</b>	A6D6

**Description:** Eukaryotic histones are basic and water soluble nuclear proteins that form hetero-octameric nucleosome particles by wrapping 146 base pairs of DNA in a left-handed super-helical turn sequentially to form chromosomal fibers. Two molecules of each of the four core histones (H2A, H2B, H3 and H4) form the octamer, which is comprised of two H2A-H2B dimers and two H3-H4 dimers, forming two nearly symmetrical halves by tertiary structure. Histones are subject to posttranslational modification by enzymes primarily on their N-terminal tails, but also in their globular domains. Human and mouse Histone H4 are subject to methylation at Lys 20, a modification that may be necessary for select DNA transactions or chromatin state transitions.

**Immunogen:** Synthetic peptide within human Histone H3 aa 20-40 (acetyl K27).

**Positive control:** PC-12 cell lysate, 293 cell lysate, MCF-7 cell lysate, NIH/3T3 cell lysate, HeLa cell lysates, HeLa, human tonsil tissue.

**Subcellular location:** Nucleus, Chromosome.

**Database links:** SwissProt P68431 Human | P84243 Human | Q16695 Human | Q6NXT2 Human | Q71DI3 Human | P68433 Mouse | Q6LED0 Rat

### Recommended Dilutions:

<b>WB</b>	1:500-1:2,000
<b>IF-Cell</b>	1:100
<b>IHC-P</b>	1:600
<b>FC</b>	1:500
<b>ChIP</b>	Use 0.5~2 µg for 25 µg of chromatin.

**Storage Buffer:** PBS (pH 7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.05% SodiumAzide.

**Storage Instruction:** Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

**Purity:** Protein G affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn


## Images



**Fig1:** Western blot analysis of Histone H3 (acetyl K27) on different lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (HA600048, 1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1:20,000 dilution was used for 1 hour at room temperature.


**Positive control:**

Lane 1: PC-12 cell lysate  
Lane 2: 293 cell lysate  
Lane 3: MCF-7 cell lysate  
Lane 4: NIH/3T3 cell lysate




**Fig2:** Western blot analysis of Histone H3 (acetyl K27) on HeLa cell lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (HA600048, 1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1:20,000 dilution was used for 1 hour at room temperature.

Lane 1: Untreated HeLa whole cell lysates  
Lane 2: HeLa treated with Trichostatin A whole cell lysates



**Fig3:** ICC staining of Histone H3 (acetyl K27) in HeLa cells (green). Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 10% negative goat serum for 15 minutes at room temperature. Cells were probed with the primary antibody (HA600048, 1/100) for 1 hour at room temperature, washed with PBS. Alexa Fluor®488 conjugate-Goat anti-Mouse IgG was used as the secondary antibody at 1/1,000 dilution. The nuclear counter stain is DAPI (blue).



**Fig4:** Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-Histone H3 (acetyl K27) antibody. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 20 minutes. The tissues were blocked in 1% BSA for 30 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA600048, 1/600) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn



**Fig5:** Flow cytometric analysis of Histone H3 (acetyl K27) was done on HeLa cells. The cells were fixed, permeabilized and stained with the primary antibody (HA600048, 1ug/ml) (red) compared with Rabbit IgG, monoclonal - Isotype Control (green). After incubation of the primary antibody at +4°C for 1 hour, the cells were stained with a Alexa Fluor®488 conjugate-Goat anti-Mouse IgG Secondary antibody at 1/1000 dilution for 30 minutes at +4°C (dark incubation). Unlabelled sample was used as a control (cells without incubation with primary antibody; black).



**Fig6:** Chromatin immunoprecipitations were performed with cross-linked chromatin from HeLa cells treated with 500ng/mL TSA for 4 hours and either Histone H3 (acetyl K27) (HA600048) or Normal Mouse IgG according to the ChIP protocol. The enriched DNA was quantified by real-time PCR using indicated primers. The amount of immunoprecipitated DNA in each sample is represented as signal relative to the total amount of input chromatin, which is equivalent to one.

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

## Background References

1. Ray-Gallet D. et al. The Histone H3 Family and Its Deposition Pathways. *Adv Exp Med Biol.* 2021
2. Francis NJ. et al. Inheritance of Histone (H3/H4): A Binary Choice? *Trends Biochem Sci.* 2021 Jan

Hangzhou Huaan Biotechnology Co., Ltd.

Orders: 0086-571-88062880

Technical: 0086-571-89986345

Service mail: support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn