

Anti-IL-1 beta Antibody [A7A10]

HA601035



Product Type:	Mouse monoclonal IgG1, primary antibodies
Species reactivity:	Human
Applications:	WB, ELISA, IF-Cell
Molecular Wt:	Predicted band size: 31 kDa
Clone number:	A7A10

Description: The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. Similarly, IL-1B has been implicated in human osteoarthritis pathogenesis. Patients with severe Coronavirus Disease 2019 (COVID-19) present elevated levels of pro-inflammatory cytokines such as IL-1B in bronchial alveolar lavage fluid samples. The lung damage induced by the Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is to a large extent, a result of the inflammatory response promoted by cytokines such as IL-1B. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2.

Immunogen: Recombinant protein within Human IL-1 beta aa 117-269 / 269.

Positive control: THP-1 treated with 80nM TPA overnight then replaced with 100ng/mL LPS for 6 hours add 300ng/mL BFA for last 3 hours cell lysate, THP-1 cells treated with 80nM TPA overnight then replaced with 100ng/mL LPS for 6 hours add 300ng/mL BFA for last 3 hours.

Subcellular location: Extracellular exosome, Secreted, Lysosome, Cytosol.

Database links: SwissProt: P01584 Human

Recommended Dilutions:

WB	1:1,000-1:2,000
ELISA	1:10,000
IF-Cell	1:100

Storage Buffer: PBS (pH7.4), 0.1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Shipped at 4°C. Store at +4°C short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20°C long term.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn


华安生物
HUABIO
www.huabio.cn

Images

Fig1: Western blot analysis of IL-1 beta on different lysates with Mouse anti-IL-1 beta antibody (HA601035) at 1/1,000 dilution.

Lane 1: THP-1 cell lysate

Lane 2: THP-1 treated with 80nM TPA overnight then replaced with 100ng/mL LPS for 6 hours add 300ng/mL BFA for last 3 hours cell lysate



Lysates/proteins at 10 µg/Lane.

Predicted band size: 31 kDa


Observed band size: 31 kDa

Exposure time: 3 minutes;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA601035) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig2: Immunocytochemistry analysis of THP-1 cells treated with 80nM TPA overnight then replaced with 100ng/mL LPS for 6 hours add 300ng/mL BFA for last 3 hours labeling IL-1 beta with Mouse anti-IL-1 beta antibody (HA601035) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Mouse anti-IL-1 beta antibody (HA601035) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 488, HA1125) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

beta Tubulin (ET1602-4, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Rabbit IgG H&L (iFluor™ 594, HA1122) were used as the secondary antibody at 1/1,000 dilution.


Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Fig3: IL-1 beta Antibody (HA601035) in indirect ELISA.

Indirect ELISA analysis of IL-1 beta was performed by coating wells of a 96-well plate with 50 μ l per well of IL-1 beta antigen diluted in carbonate/bicarbonate buffer, at a concentration of 1 μ g/mL overnight at 4°C. Wells of the plate were washed, blocked with StartingBlock blocking buffer, and incubated with 50 μ l per well of a mouse IL-1 beta monoclonal antibody starting at a concentration of 20 μ g/mL and serially diluting it to a concentration of 1.28 ng/mL for 2 hours at room temperature. The plate was washed and incubated with 50 μ l per well of an HRP-conjugated goat anti-mouse IgG secondary antibody at a dilution of 1:10,000 for one hour at room temperature. Detection was performed using an Ultra TMB Substrate for 5 minutes at room temperature in the dark. The reaction was stopped with sulfuric acid and absorbances were read on a spectrophotometer at 450 nm.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Zou L. et. al. HO-1 induced autophagy protects against IL-1 beta-mediated apoptosis in human nucleus pulposus cells by inhibiting NF-kappaB. *Aging* (Albany NY). 2020 Feb
2. Libby P. Interleukin-1 Beta as a Target for Atherosclerosis Therapy: Biological Basis of CANTOS and Beyond. *J Am Coll Cardiol*. 2017 Oct

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn