

# Anti-GNAS Antibody [A8F12]

HA601074



<b>Product Type:</b>	Mouse monoclonal IgG2a, primary antibodies
<b>Species reactivity:</b>	Human, Mouse, Rat
<b>Applications:</b>	WB, IHC-P, IF-Cell
<b>Molecular Wt:</b>	Predicted band size: 46 kDa
<b>Clone number:</b>	A8F12

**Description:** This locus has a highly complex imprinted expression pattern. It gives rise to maternally, paternally, and biallelically expressed transcripts that are derived from four alternative promoters and 5' exons. Some transcripts contain a differentially methylated region (DMR) at their 5' exons, and this DMR is commonly found in imprinted genes and correlates with transcript expression. An antisense transcript is produced from an overlapping locus on the opposite strand. One of the transcripts produced from this locus, and the antisense transcript, are paternally expressed noncoding RNAs, and may regulate imprinting in this region. In addition, one of the transcripts contains a second overlapping ORF, which encodes a structurally unrelated protein - Alex. Alternative splicing of downstream exons is also observed, which results in different forms of the stimulatory G-protein alpha subunit, a key element of the classical signal transduction pathway linking receptor-ligand interactions with the activation of adenylyl cyclase and a variety of cellular responses. Multiple transcript variants encoding different isoforms have been found for this gene. Mutations in this gene result in pseudohypoparathyroidism type 1a, pseudohypoparathyroidism type 1b, Albright hereditary osteodystrophy, pseudopseudohypoparathyroidism, McCune-Albright syndrome, progressive osseous heteroplasia, polyostotic fibrous dysplasia of bone, and some pituitary tumors.

**Immunogen:** Recombinant protein within human GNAS aa 2-251.

**Positive control:** HepG2 cell lysate, NIH/3T3 cell lysate, human brain tissue lysate, rat pancreas tissue, human pancreas tissue, MCF-7, human breast carcinoma tissue.

**Subcellular location:** Cell membrane.

**Database links:** SwissProt: P63092 Human | P84996 Human | Q5JWF2 Human | P63094 Mouse | Q6R0H7 Mouse | P63095 Rat | Q63803 Rat

**Recommended Dilutions:**

<b>WB</b>	1:1,000
<b>IHC-P</b>	1:200-1:500
<b>IF-Cell</b>	1:200

**Storage Buffer:** PBS (pH7.4), 0.1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

**Storage Instruction:** Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

**Purity:** Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

## Images



**Fig1:** Western blot analysis of GNAS on different lysates with Mouse anti-GNAS antibody (HA601074) at 1/1,000 dilution.

Lane 1: HepG2 cell lysate (10 µg/Lane)

Lane 2: NIH/3T3 cell lysate (10 µg/Lane)

Lane 3: Human brain tissue lysate (20 µg/Lane)


Predicted band size: 46 kDa

Observed band size: 46 kDa

Exposure time: 2 minutes;


4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody (HA601074) at 1/1,000 dilution was used in 5% NFDN/TBST at room temperature for 2 hours. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1:100,000 dilution was used for 1 hour at room temperature.



**Fig2:** Immunohistochemical analysis of paraffin-embedded rat pancreas tissue with Mouse anti-GNAS antibody (HA601074) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA601074) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig3:** Immunohistochemical analysis of paraffin-embedded human pancreas tissue with Mouse anti-GNAS antibody (HA601074) at 1/500 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA601074) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn


华安生物  
HUABIO  
www.huabio.cn



**Fig4:** Immunocytochemistry analysis of MCF-7 cells labeling GNAS with Mouse anti-GNAS antibody (HA601074) at 1/200 dilution.

Cells were fixed in 4% paraformaldehyde for 30 minutes, permeabilized with 0.05% Triton X-100 in PBS for 20 minutes, and then blocked with 2% negative goat serum for 30 minutes at room temperature. Cells were then incubated with Mouse anti-GNAS antibody (HA601074) at 1/200 dilution in 2% negative goat serum overnight at 4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 488, HA1125) was used as the secondary antibody at 1/1,000 dilution. Nuclear DNA was labelled in blue with DAPI.

beta Tubulin (ET1602-4, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Rabbit IgG H&L (iFluor™ 647, HA1123) were used as the secondary antibody at 1/1,000 dilution.



**Fig5:** Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue with Rabbit anti-GNAS antibody (HA601074) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA601074) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

### Background References

1. Thiele S., de Sanctis L., Werner R., Grotzinger J., Aydin C., Juppner H., Bastepe M., Hiort O. Functional characterization of GNAS mutations found in patients with pseudohypoparathyroidism type 1c defines a new subgroup of pseudohypoparathyroidism affecting selectively Gsalpha-receptor interaction. *Hum. Mutat.* 32:653-660(2011)
2. Brand C.S., Sadana R., Malik S., Smrcka A.V., Dessauer C.W. Adenylyl cyclase 5 regulation by Gbetagamma involves isoform specific use of multiple interaction sites. *Mol. Pharmacol.* 88:758-767(2015)

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn