

# Anti-S100 beta Antibody [PD00-11]

HA601098



<b>Product Type:</b>	Recombinant Mouse monoclonal IgG, primary antibodies
<b>Species reactivity:</b>	Human
<b>Applications:</b>	IHC-P, WB
<b>Molecular Wt:</b>	Predicted band size: 11 kDa
<b>Clone number:</b>	PD00-11

**Description:** S100 calcium binding protein B (S100 beta) is a member of the multifunctional S100 family of proteins. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21; however, this gene is located at 21q22.3. S100 beta acts as a stimulator of proliferation and migration and as an inhibitor of apoptosis and differentiation in many cell types including astrocytes, Schwann cells, chondrocytes, adipocytes, certain neuronal populations, melanocytes, Langerhans cells, histiocytes, epithelial, and myoepithelial cells. This protein may function in Neurite extension, proliferation of melanoma cells, stimulation of Ca<sup>2+</sup> fluxes, inhibition of PKC-mediated phosphorylation, astrocytosis and axonal proliferation, and inhibition of microtubule assembly. Chromosomal rearrangements and altered expression of this gene have been implicated in several neurological, neoplastic, and other types of diseases, including Alzheimer's disease, Down's syndrome, epilepsy, amyotrophic lateral sclerosis, melanoma, and type I diabetes. S100 beta is also expressed in neoplasms derived from these cell types, making it a useful marker for the identification of melanoma and various nervous system tumors. Although ubiquitous, S100 beta has proven to be a sensitive marker for malignant melanoma, including desmoplastic and metastatic variants.

<b>Immunogen:</b>	Full length corresponding to Human S100 beta.
<b>Positive control:</b>	SK-MEL-28 cell lysates, human malignant melanoma tissue, human meningioma tissue, human tonsil tissue.
<b>Subcellular location:</b>	Cytoplasm, Nucleus.
<b>Database links:</b>	SwissProt P04271 Human
<b>Recommended Dilutions:</b>	
IHC-P	1:5,000
WB	1:1,000
<b>Storage Buffer:</b>	PBS (pH7.4), 0.1% BSA, 40% Glycerol. Preservative: 0.05% SodiumAzide.
<b>Storage Instruction:</b>	Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.
<b>Purity:</b>	Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

**Fig1:** Western blot analysis of S100 beta on SK-MEL-28 cell lysates with Rabbit anti-S100 beta antibody (HA601098) at 1/1,000 dilution.

Lysates/proteins at 10 µg/Lane.



Predicted band size: 11 kDa

Observed band size: 11 kDa

Exposure time: 1 minute; ECL: K1801;


4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFD/MTBST for 1 hour at room temperature. The primary antibody (HA601098) at 1/1,000 dilution was used in 5% NFD/MTBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.




**Fig2:** Immunohistochemical analysis of paraffin-embedded human malignant melanoma tissue with Mouse anti-S100 beta antibody (HA601098) at 1/5,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA601098) at 1/5,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig3:** Immunohistochemical analysis of paraffin-embedded human meningioma tissue with Mouse anti-S100 beta antibody (HA601098) at 1/5,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA601098) at 1/5,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig4:** Immunohistochemical analysis of paraffin-embedded human tonsil tissue with Mouse anti-S100 beta antibody (HA601098) at 1/5,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA601098) at 1/5,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

### Background References

1. Miyamoto Y et al. Involvement of the Tyro3 receptor and its intracellular partner Fyn signaling in Schwann cell myelination. *Mol Biol Cell* 26:3489-503 (2015).
2. Gondo A et al. Sustained Down-regulation of -Dystroglycan and Associated Dysfunctions of Astrocytic Endfeet in Epileptic Cerebral Cortex. *J Biol Chem* 289:30279-88 (2014).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn