

Anti-HSP90 Beta Antibody [2-1-G3-R]

HA601176



Product Type:	Recombinant Mouse monoclonal IgG1, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IHC-P, IF-Cell
Molecular Wt:	Predicted band size: 83.3 kDa
Clone number:	2-1-G3-R

Description: Molecular chaperone that promotes the maturation, structural maintenance and proper regulation of specific target proteins involved for instance in cell cycle control and signal transduction. Undergoes a functional cycle linked to its ATPase activity. This cycle probably induces conformational changes in the client proteins, thereby causing their activation. Interacts dynamically with various co-chaperones that modulate its substrate recognition, ATPase cycle and chaperone function. Engages with a range of client protein classes via its interaction with various co-chaperone proteins or complexes, that act as adapters, simultaneously able to interact with the specific client and the central chaperone itself. Recruitment of ATP and co-chaperone followed by client protein forms a functional chaperone. After the completion of the chaperoning process, properly folded client protein and co-chaperone leave HSP90 in an ADP-bound partially open conformation and finally, ADP is released from HSP90 which acquires an open conformation for the next cycle. Apart from its chaperone activity, it also plays a role in the regulation of the transcription machinery. HSP90 and its co-chaperones modulate transcription at least at three different levels. They first alter the steady-state levels of certain transcription factors in response to various physiological cues. Second, they modulate the activity of certain epigenetic modifiers, such as histone deacetylases or DNA methyl transferases, and thereby respond to the change in the environment. Third, they participate in the eviction of histones from the promoter region of certain genes and thereby turn on gene expression.

Immunogen:	Synthetic peptide within Human HSP90 Beta aa 311-360 / 724.
Positive control:	HeLa cell lysate, HepG2 cell lysate, A549 cell lysate, K-562 cell lysate, Jurkat cell lysate, NIH/3T3 cell lysate, C6 cell lysate, PC-12 cell lysate, mouse brain tissue lysate, mouse heart tissue lysate, rat brain tissue lysate, human brain tissue, rat brain tissue, HeLa.
Subcellular location:	Cytoplasm
Database links:	SwissProt: P08238 Human P11499 Mouse P34058 Rat
Recommended Dilutions:	
WB	1:1,000-1:2,000
IHC-P	1:5,000
IF-Cell	1:100
Storage Buffer:	PBS (pH7.4), 0.1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.
Storage Instruction:	Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.
Purity:	Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images

Fig1: Western blot analysis of HSP90 Beta on different lysates with Mouse anti-HSP90 Beta antibody (HA601176) at 1/1,000 dilution.



Lane 1: HeLa cell lysate (20 µg/Lane)
 Lane 2: HepG2 cell lysate (20 µg/Lane)
 Lane 3: A549 cell lysate (20 µg/Lane)
 Lane 4: K-562 cell lysate (20 µg/Lane)
 Lane 5: Jurkat cell lysate (20 µg/Lane)
 Lane 6: NIH/3T3 cell lysate (20 µg/Lane)
 Lane 7: C6 cell lysate (20 µg/Lane)
 Lane 8: PC-12 cell lysate (20 µg/Lane)
 Lane 9: Mouse brain tissue lysate (40 µg/Lane)
 Lane 10: Mouse heart tissue lysate (40 µg/Lane)
 Lane 11: Rat brain tissue lysate (40 µg/Lane)

Predicted band size: 83 kDa

Observed band size: 83 kDa

Exposure time: 53 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDm/TBST for 1 hour at room temperature. The primary antibody (HA601176) at 1/1,000 dilution was used in 5% NFDm/TBST at room temperature for 2 hours. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/100,000 dilution was used for 1 hour at room temperature.




Fig2: Immunohistochemical analysis of paraffin-embedded human brain tissue with Mouse anti-HSP90 Beta antibody (HA601176) at 1/5,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA601176) at 1/5,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn





Fig3: Immunohistochemical analysis of paraffin-embedded rat brain tissue with Mouse anti-HSP90 Beta antibody (HA601176) at 1/5,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA601176) at 1/5,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Fig4: Immunocytochemistry analysis of HeLa cells labeling HSP90 Beta with Mouse anti-HSP90 Beta antibody (HA601176) at 1/100 dilution.



Cells were fixed in 100% precooled methanol for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Mouse anti-HSP90 Beta antibody (HA601176) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 488, HA1125) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

beta Tubulin (ET1602-4, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Rabbit IgG H&L (iFluor™ 594, HA1122) were used as the secondary antibody at 1/1,000 dilution.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Chadli A., Graham J.D., Abel M.G., Jackson T.A., Gordon D.F., Wood W.M., Felts S.J., Horwitz K.B., Toft D. GCUNC-45 is a novel regulator for the progesterone receptor/hsp90 chaperoning pathway. *Mol. Cell. Biol.* 26:1722-1730 (2006)
2. Verma S., Goyal S., Jamal S., Singh A., Grover A. Hsp90: Friends, clients and natural foes. *Biochimie* 127:227-240 (2016)

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn