

iFluor™ 647 Conjugated Anti-Cytokeratin 7 Antibody [ST50-05]

HA720144F



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Mouse
Applications:	IF-Cell, IF-Tissue
Molecular Wt:	Predicted band size: 51 kDa
Clone number:	ST50-05

Description: Cytokeratins comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue, where they constitute up to 85% of mature keratinocytes in the vertebrate epidermis. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. The α -helical coiled-coil dimers associate laterally end-to-end to form 10 nm diameter filaments. Cytokeratins are useful markers of tissue differentiation and, in addition, they aid in the characterization of malignant tumors. Cytokeratin 7 (also known as sarcolectin) agglutinates normal and transformed cells with a high affinity for simple sugars. Cytokeratin 7 also inhibits the synthesis of interferon-dependent secondary proteins thus reversing the antiviral effect of interferon induction and restoring cells to their status ad primum. In normal and transformed cells, Cytokeratin 7 localizes to the membrane.

Conjugate: iFluor™ 647, Ex: 656nm; Em: 670nm.

Immunogen: Synthetic peptide within Human Cytokeratin 7 aa 18-67.

Positive control: SK-Br-3, human liver tissue, human breast tissue.

Subcellular location: Cytoplasm.

Database links: SwissProt: P08729 Human | Q9DCV7 Mouse

Recommended Dilutions:

IF-Cell	1:100
IF-Tissue	1:50-1:200

Storage Buffer: Preservative: 0.02% Sodium azide Constituents: 30% Glycerol, 1% BSA, 68.98% PBS.

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images




Fig1: Immunocytochemistry analysis of SK-Br-3 cells labeling Cytokeratin 7 with Rabbit anti-Cytokeratin 7 antibody (HA720144F) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 15 minutes, permeabilized with 0.1% Triton X-100 in PBS for 15 minutes, and then blocked with 2% normal goat serum for 1 hour at 37 °C. Cells were then incubated with Rabbit anti-Cytokeratin 7 antibody (HA720144F, red) at 1/100 dilution in 2% normal goat serum overnight at 4 °C. Nuclear DNA was labelled in blue with DAPI.

Beta III Tubulin (M0805-8, green) was stained at 1/200 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 488, HA1125) was used as the secondary antibody at 1/800 dilution.




Fig2: Immunofluorescence analysis of paraffin-embedded human liver tissue labeling Cytokeratin 7 (HA720144F) and Cytokeratin 8 (HA720132F).

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 10% negative goat serum for 1 hour at room temperature, washed with PBS. And then probed with the primary antibodies Cytokeratin 7 (HA720144F, red) at 1/200 dilution and Cytokeratin 8 (HA720132F, green) at 1/200 dilution overnight at 4 °C, washed with PBS.

DAPI was used as nuclear counterstain.




Fig3: Immunofluorescence analysis of paraffin-embedded human breast tissue labeling Cytokeratin 7 (HA720144F), Cytokeratin 8 (HA720132F) and Vimentin (EM0401).

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 10% negative goat serum for 1 hour at room temperature, washed with PBS. And then probed with the primary antibodies Cytokeratin 7 (HA720144F, red) at 1/50 dilution, Cytokeratin 8 (HA720132F, green) at 1/200 dilution and Vimentin (EM0401, yellow) at 1/1,000 dilution overnight at 4 °C, washed with PBS.

Alexa Fluor® 555 conjugate-Goat anti-Mouse IgG was used as the secondary antibody at 1/1,000 dilution. DAPI was used as nuclear counterstain.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Hrudka J. et al. Cytokeratin 7 expression as a predictor of an unfavorable prognosis in colorectal carcinoma. *Sci Rep.* 2021 Sep
2. Statz E. et al. Cytokeratin 7, GATA3, and SOX-10 is a Comprehensive Panel in Diagnosing Triple Negative Breast Cancer Brain Metastases. *Int J Surg Pathol.* 2021 Aug

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUAABIO
www.huabio.cn