Anti-IL-1 beta Antibody [PS00-82] HA721279

Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human
Applications:	ELISA(Det)
Molecular Wt:	Predicted band size: 31 kDa
Clone number:	PS00-82
Description:	Interleukin-1 beta (IL-1 β) also known as leukocytic pyrogen, leukocytic endogenous mediator, mononuclear cell factor, lymphocyte activating factor and other names, is a cytokine protein that in humans is encoded by the IL1B gene. There are two genes for interleukin-1 (IL-1): IL-1 alpha and IL-1 beta (this gene). IL-1 β precursor is cleaved by cytosolic caspase 1 (interleukin 1 beta convertase) to form mature IL-1 β . Increased production of IL-1 β causes a number of different autoinflammatory syndromes, most notably the monogenic conditions referred to as Cryopyrin-Associated Periodic Syndromes (CAPS), due to mutations in the inflammasome receptor NLRP3 which triggers processing of IL-1B. Intestinal dysbiosis has been observed to induce osteomyelitis through a IL-1 β dependent manner. The presence of IL-1 β has been also found in patients with multiple sclerosis (a chronic autoimmune disease of the central nervous system). However, it is not known exactly which cells produce IL-1 β . Treatment of multiple sclerosis with glatiramer acetate or natalizumab has also been shown to reduce the presence of IL-1 β or its receptor.[18]
lmmunogen:	Recombinant protein within full length human IL-1 beta.
Positive control:	Recombinant IL-1 beta protein
Subcellular location:	Cytoplasm, Lysosome, Secreted
Database links:	SwissProt: P01584 Human
Recommended Dilutions: ELISA (Det)	Use at an assay dependent concentration. Can be paired for Sandwich ELISA with Rabbit monoclonal [PS01-39] to IL-1 beta (Capture) (HA721278).
Storage Buffer:	PBS (pH7.4).
Storage Instruction:	Store at +4 $^\circ\!\mathrm{C}$ after thawing. Aliquot store at -20 $^\circ\!\mathrm{C}$. Avoid repeated freeze / thaw cycles.
Purity:	Protein A affinity purified.

Hangzhou HuaAn Biotechnology Co.,Ltd.Orders: 0086-571-88062880Technical:0086-571-89986345

Service mail: support@huabio.cn





11.

Applications: WB=Western blot IP=Immunoprecipitation IHC=Immunohistochemistry IF=Immunofluorescence FC=Flow cytometry

HA721279 - Page 2

Images





beta protein (Con. pg/ml)

Standard curve of IL-1 beta matched pair antibodies

Fig1: The binding activity of IL-1 beta (HA721279) with recombinant IL-1 beta protein.

Immobilized recombinant IL-1 beta protein at 1 µg/ml overnight at 4°C. Then blocked with 1xTBS/1%BSA for 1 hour at 37°C, and incubated with the primary antibody (HA721279) for 45min at 37°C. Then the plate was washed and incubated with 50 µl per well of Goat anti-Rabbit IgG-HRP for 0.5 hour at 37°C. Detection was performed using an Ultra TMB Substrate for 10 minutes at room temperature in the dark. The reaction was stopped with sulfuric acid and absorbances were read on a spectrophotometer at 450 nm.

Fig2: Standard curve of IL-1 beta matched pair antibodies:

Sandwich ELISA analysis of IL-1 beta matched pair antibodies Elisa assay was performed by coating wells of a 96-well plate with 50 µl per well of capture antibody HA721278 [PS01-39] diluted in carbonate/bicarbonate buffer, at a concentration of 4 µg/mL overnight at 4°C. Wells of the plate were washed, blocked with 150 µl 1% BSA/PBST blocking buffer, and incubated with serial diluted recombinant IL-1 beta protein starting from 2000 pg/ml to 31.25 pg/ml for 1 hour at 37°C. The plate was washed and incubated with 50 µl per well of detect antibody [PS00-82] (Biotin, 1:2,000) for 1 hour at 37°C. Then the plate was washed and incubated with 50 µl per well of Streptavidin-HRP for 0.5 hour at 37°C. Detection was performed using an Ultra TMB Substrate for 10 minutes at room temperature in the dark. The reaction was stopped with sulfuric acid and absorbances were read on a spectrophotometer at 450 nm.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

- 1. Milosevic V, et al. (January 2020). "Wnt/IL-1β/IL-8 autocrine circuitries control chemoresistance in mesothelioma initiating cells by inducing ABCB5". Int. J. Cancer. 146 (1): 192–207.
- Roy D, Sarkar S, Felty Q (January 2006). "Levels of IL-1 beta control stimulatory/inhibitory growth of cancer cells". Frontiers in Bioscience. 11: 889–98.

Hangzhou HuaAn Biotechnology Co.,Ltd. Orders: 0086-571-88062880 Technical:008

Technical:0086-571-89986345

Service mail: support@huabio.cn



www.huabio.cn

Applications: WB=Western blot IP=Immunoprecipitation IHC=Immunohistochemistry IF=Immunofluorescence FC=Flow cytometry