

# Anti-Phospho-Histone H3 (T3) Antibody [JE42-48]

## HA721796



<b>Product Type:</b>	Recombinant Rabbit monoclonal IgG, primary antibodies
<b>Species reactivity:</b>	Human, Mouse, Rat
<b>Applications:</b>	WB, IHC-P
<b>Molecular Wt:</b>	Predicted band size: 15 kDa
<b>Clone number:</b>	JE42-48

<b>Description:</b>	Variant histone H3 which replaces conventional H3 in a wide range of nucleosomes in active genes. Constitutes the predominant form of histone H3 in non-dividing cells and is incorporated into chromatin independently of DNA synthesis. Deposited at sites of nucleosomal displacement throughout transcribed genes, suggesting that it represents an epigenetic imprint of transcriptionally active chromatin. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
<b>Immunogen:</b>	Synthetic phosphopeptide corresponding to residues surrounding Thr3 of Human Histone H3.
<b>Positive control:</b>	HeLa treated with 20% FBS and 100nM Calyculin A for 30 minutes cell lysate, NIH/3T3 treated with 20% PBS and 100nM Calyculin A for 30 minutes cell lysate, mouse breast tissue, mouse skin tissue.
<b>Subcellular location:</b>	Nucleus, Chromosome.
<b>Database links:</b>	SwissProt: P68431 Human   P68433 Mouse   Q6LED0 Rat
<b>Recommended Dilutions:</b>	
<b>WB</b>	1:1,000
<b>IHC-P</b>	1:200-1:1,000
<b>Storage Buffer:</b>	1*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.
<b>Storage Instruction:</b>	Shipped at 4℃. Store at +4℃ short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20℃ long term.
<b>Purity:</b>	Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders: 0086-571-88062880


Technical: 0086-571-89986345

Service mail: support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn

Applications: WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

## Images



**Fig1:** Western blot analysis of Phospho-Histone H3 (T3) on different lysates with Rabbit anti-Phospho-Histone H3 (T3) antibody (HA721796) at 1/1,000 dilution.

Lane 1: HeLa cell lysate

Lane 2: HeLa treated with 20% FBS and 100nM Calyculin A for 30 minutes cell lysate

Lane 3: NIH/3T3 cell lysate

Lane 4: NIH/3T3 treated with 20% PBS and 100nM Calyculin A for 30 minutes cell lysate

Lysates/proteins at 20 µg/Lane.


Predicted band size: 15 kDa

Observed band size: 15 kDa

Exposure time: 25 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA721796) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.



**Fig2:** Immunohistochemical analysis of paraffin-embedded mouse breast tissue with Rabbit anti-Phospho-Histone H3 (T3) antibody (HA721796) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA721796) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn



**Fig3:** Immunohistochemical analysis of paraffin-embedded mouse skin tissue with Rabbit anti-Phospho-Histone H3 (T3) antibody (HA721796) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA721796) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

### Background References

1. Strobino M. et. al. Loss of histone H3.3 results in DNA replication defects and altered origin dynamics in *C. elegans*. *Genome Res.* 2020 Dec
2. Ince AT. et. al. The relationship of Serum Histone H3.3 and H4 with chronic Hepatitis B. *J Pak Med Assoc.* 2020 Sep

Hangzhou Huaan Biotechnology Co., Ltd.

Orders: 0086-571-88062880

Technical: 0086-571-89986345

Service mail: support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn

Applications: WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation