# Anti-Phospho-Tau (S324) Antibody [JE30-73] HA721798

Product Type: Species reactivity: Applications: Molecular Wt: Clone number:	Recombinant Rabbit monoclonal IgG, primary antibodies Human, Mouse, Rat WB Predicted band size: 79 kDa JE30-73
Description:	The tau proteins (abbreviated from tubulin associated unit are a group of six highly soluble protein isoforms produced by alternative splicing from the gene MAPT (microtubule-associated protein tau). They have roles primarily in maintaining the stability of microtubules in axons and are abundant in the neurons of the central nervous system (CNS), where the cerebral cortex has the highest abundance. They are less common elsewhere but are also expressed at very low levels in CNS astrocytes and oligodendrocytes. Pathologies and dementias of the nervous system such as Alzheimer's disease and Parkinson's disease are associated with tau proteins that have become hyperphosphorylated insoluble aggregates called neurofibrillary tangles. The tau proteins were identified in 1975 as heat-stable proteins essential for microtubule assembly, and since then they have been characterized as intrinsically disordered proteins.
lmmunogen:	Synthetic phosphopeptide corresponding to residues surrounding S324 of human Tau protein.
Positive control:	SH-SY5Y treated with 200nM Calyculin A and $1\mu M$ Okadaic Acid for 1 hour cell lysate.
Subcellular location:	Cytoplasm, cytosol, Cell membrane, cytoskeleton, Cell projection, axon, dendrite, Secreted.
Database links:	SwissProt: P10636-8 Human   P10637 Mouse   P19332 Rat
Recommended Dilutions: WB	1:1,000
Storage Buffer:	1*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.
Storage Instruction:	Store at +4 $^\circ\!\mathrm{C}$ after thawing. Aliquot store at -20 $^\circ\!\mathrm{C}$ . Avoid repeated freeze / thaw cycles.
Purity:	Protein A affinity purified.

## Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn



11.

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

#### Images

**Fig1:** Western blot analysis of Phospho-Tau (S324) on different lysates with Rabbit anti-Phospho-Tau (S324) antibody (HA721798) at 1/1,000 dilution.

Lane 1: SH-SY5Y cell lysate Lane 2: SH-SY5Y treated with 200nM Calyculin A and 1µM Okadaic Acid for 1 hour cell lysate

Lysates/proteins at 20 µg/Lane.

Predicted band size: 79 kDa Observed band size: 35-130 kDa

Exposure time: 1 minute;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA721798) at 1/1,000 dilution was used in 5% NFDM/TBST at  $4^{\circ}$ C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

#### **Background References**

- 1. Karikari TK et al. Blood phospho-tau in Alzheimer disease: analysis, interpretation, and clinical utility. Nat Rev Neurol. 2022 Jul
- 2. Palmqvist S et al. Prediction of future Alzheimer's disease dementia using plasma phospho-tau combined with other accessible measures. Nat Med. 2021 Jun

### Hangzhou Huaan Biotechnology Co., Ltd.



Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

