Anti-beta Actin Antibody [PSH03-63] HA722023

Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Mouse, Rat, Rabbit, Goat, Zebrafish, Dog, Monkey, Pig, Chicken, Cow, Hamster, Green monkey
Applications:	WB, IF-Cell, FC, IP
Molecular Wt:	Predicted band size: 42 kDa
Clone number:	PSH03-63
Description:	Actins are highly conserved proteins involved in cell motility, structure and integrity. Actin has been found to be expressed in at least six isomeric forms. It is expressed in heart and skeletal striated muscle tissue, and in certain smooth muscle tissues, regulating contractile potentials for these cells. It is also expressed in the cytoplasm of non-muscle cells, functioning to control cell structure and motility. Beta actin is usually used as a loading control, for among others, the integrity of cells, protein degradation, in Western Blotting.
lmmunogen:	Synthetic peptide within N-terminal residues of β -Actin.
Positive control:	HeLa cell lysate, HepG2 cell lysate, MCF7 cell lysate, A431 cell lysate, Jurkat cell lysate, HEK-293 cell lysate, RAW264.7 cell lysate, C2C12 cell lysate, PC-12 cell lysate, mouse testis tissue lysate, mouse spleen tissue lysate, rat testis tissue lysate, rat kidney tissue lysate, HeLa, NIH/3T3, C6.
Subcellular location:	Cytoskeleton.
Database links:	SwissProt: P60709 Human P60710 Mouse P60711 Rat
Recommended Dilutions: WB IF-Cell FC IP	1:20,000-1:100,000 1:100-1:250 1:1,000 1-2µg/sample
Storage Buffer:	PBS (pH7.4), 0.1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.
Storage Instruction:	Shipped at 4 $^\circ\!\!\mathbb{C}$. Store at +4 $^\circ\!\!\mathbb{C}$ short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20 $^\circ\!\!\mathbb{C}$ long term.
Purity:	Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

45 Service mail:support@huabio.cn



11.

Images



Fig1: Western blot analysis of beta Actin on different lysates with Rabbit anti-beta Actin antibody (HA722023) at 1/20,000 dilution.

- Lane 1: HeLa cell Iysate (10 µg/Lane) Lane 2: HepG2 cell Iysate (10 µg/Lane) Lane 3: MCF7 cell Iysate (10 µg/Lane) Lane 4: A431 cell Iysate (10 µg/Lane) Lane 5: Jurkat cell Iysate (10 µg/Lane) Lane 6: HEK-293 cell Iysate (10 µg/Lane) Lane 7: RAW264.7 cell Iysate (10 µg/Lane) Lane 8: C2C12 cell Iysate (10 µg/Lane) Lane 9: PC-12 cell Iysate (10 µg/Lane) Lane 10: Mouse testis tissue Iysate (10 µg/Lane) Lane 11: Mouse spleen tissue Iysate (10 µg/Lane) Lane 12: Rat testis tissue Iysate (10 µg/Lane)
- Lane 13: Rat kidney tissue lysate (10 µg/Lane)

Predicted band size: 42 kDa Observed band size: 42 kDa

Exposure time: 3 seconds; 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA722023) at 1/20,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig2: Western blot analysis of beta Actin on HepG2 cell lysates with Rabbit anti-beta Actin antibody (HA722023) at different dilutions.

Lysates/proteins at 20 µg/Lane.

Predicted band size: 42 kDa Observed band size: 42 kDa

Exposure time: 30 seconds; 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA722023) at different dilutions was used in 5% NFDM/TBST at 4° C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Hangzhou Huaan Biotechnology Co., Ltd.

1:320,000

Beta Actin

12kDa

1.160,000

A0,000,000

,20,000

kDa 250-

150-

100 72

55

45

35 25

14

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn





Fig3: Immunocytochemistry analysis of HeLa cells labeling beta Actin with Rabbit anti-beta Actin antibody (HA722023) at 1/250 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-beta Actin antibody (HA722023) at 1/250 dilution in 1% BSA in PBST overnight at 4 $^{\circ}$ C. Goat Anti-Rabbit IgG H&L (iFluorTM 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor ™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.



Fig4: Immunocytochemistry analysis of NIH/3T3 cells labeling beta Actin with Rabbit anti-beta Actin antibody (HA722023) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-beta Actin antibody (HA722023) at 1/100 dilution in 1% BSA in PBST overnight at 4 $^{\circ}$ C. Goat Anti-Rabbit IgG H&L (iFluorTM 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4 $^{\circ}$ C. Goat Anti-Mouse IgG H&L (iFluor TM 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn





Fig5: Immunocytochemistry analysis of L6 cells labeling beta Actin with Rabbit anti-beta Actin antibody (HA722023) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-beta Actin antibody (HA722023) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at $+4^{\circ}$ C. Goat Anti-Mouse IgG H&L (iFluor = 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.



Fig6: Flow cytometric analysis of HeLa cells labeling beta Actin.

Cells were fixed and permeabilized. Then stained with the primary antibody (HA722023, 1µg/mL) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4 °C for an hour, the cells were stained with a iFluor TM 488 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1121) at 1/1,000 dilution for 30 minutes at +4 °C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).



Fig7: Flow cytometric analysis of NIH/3T3 cells labeling beta Actin.

Cells were fixed and permeabilized. Then stained with the primary antibody (HA722023, 1µg/mL) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4°C for an hour, the cells were stained with a iFluor[™] 488 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1121) at 1/1,000 dilution for 30 minutes at +4°C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn



Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

Orders:0086-571-88062880

 Ponte P., Ng S.Y., Engel J., Gunning P., Kedes L."Evolutionary conservation in the untranslated regions of actin mRNAs: DNA sequence of a human beta-actin cDNA." Nucleic Acids Res. 12:1687-1696(1984) Ohmori H., Toyama S., Toyama S."Direct proof that the primary site of action of cytochalasin on cell motility processes is actin."J. Cell Biol. 116:933-941(1992)

Hangzhou Huaan Biotechnology Co., Ltd.



Service mail:support@huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

Technical:0086-571-89986345