

Anti-Histone H3 (acetyl K4) Antibody [PSH03-71]

HA722035



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IF-Cell, ChIP, Dot Blot
Molecular Wt:	Predicted band size: 15 kDa
Clone number:	PSH03-71

Description: In eukaryotes, DNA is wrapped around histone octamers to form the basic unit of chromatin structure. The octamer is composed of histones H2A, H2B, H3 and H4, and it associates with approximately 200 base pairs of DNA to form the nucleosome. The association of DNA with histones results in dense packing of chromatin, which restricts proteins involved in gene transcription from binding to DNA. p300 preferentially acetylates Histone H3 at lysines 14 and 18 and Histone H4 at lysines 5 and 8. PCAF in its native form, primarily acetylates Histone H3 at lysine 14 to a monoacetylated form, and less efficiently acetylates Histone H4 at lysine 8. Histone H4 may also be acetylated at lysines 12 and 16, and the involvement of acetylated H4 with Histones H2A, H2B and H3 suggests that acetylated histones may be involved in dynamic chromatin remodeling.

Immunogen: Synthetic peptide within human Histone H3 (acetyl K4) aa 1-50.

Positive control: HeLa cell lysate, HeLa treated with 500ng/mL TSA for 4 hours cell lysate, NIH/3T3 treated with 400nM TSA for 18 hours cell lysate, HeLa cells treated with 500ng/mL TSA for 4 hours, C6 cells treated with 500ng/mL TSA for 6 hours.

Subcellular location: Nucleus, Chromosome.

Database links: SwissProt: P68431 human | P84243 human | Q16695 human | Q6NXT2 human | Q71D13 human | P68433 mouse | P84228 mouse | Q6LED0 rat

Recommended Dilutions:

WB	1:1,000
IF-Cell	1:500-1:1,000
ChIP	Use 0.5~2 µg for 25 µg of chromatin.
Dot Blot	1:1,000

Storage Buffer: PBS (pH7.4), 0.1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images


Fig1: Western blot analysis of Histone H3 (acetyl K4) on different lysates with Rabbit anti-Histone H3 (acetyl K4) antibody (HA722035) at 1/1,000 dilution and competitor's antibody at 1/1,000 dilution.

Lane 1: HeLa cell lysate

Lane 2: HeLa treated with 500ng/mL TSA for 4 hours cell lysate

Lane 3: NIH/3T3 cell lysate

Lane 4: NIH/3T3 treated with 400nM TSA for 18 hours cell lysate



Lysates/proteins at 20 µg/Lane.

Predicted band size: 15 kDa


Observed band size: 15 kDa

Exposure time: 1 minute 22 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDm/TBST for 1 hour at room temperature. The primary antibody (HA722035) at 1/1,000 dilution and competitor's antibody at 1/1,000 dilution were used in 5% NFDm/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig2: Immunocytochemistry analysis of HeLa cells treated with or without 500ng/mL TSA for 4 hours labeling Histone H3 (acetyl K4) with Rabbit anti-Histone H3 (acetyl K4) antibody (HA722035) at 1/500 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Histone H3 (acetyl K4) antibody (HA722035) at 1/500 dilution in 1% BSA in PBST overnight at 4°C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn




Fig3: Immunocytochemistry analysis of C6 cells treated with or without 500ng/mL TSA for 6 hours labeling Histone H3 (acetyl K4) with Rabbit anti-Histone H3 (acetyl K4) antibody (HA722035) at 1/1,000 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Histone H3 (acetyl K4) antibody (HA722035) at 1/1,000 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

ChIP: Acetyl-Histone H3 (Lys4) HA722035




Fig4: Chromatin immunoprecipitations were performed with cross-linked chromatin from HeLa cells treated 500ng/mL TSA for 4 hours and either Histone H3 (acetyl K4) (HA722035) or Normal Rabbit IgG according to the ChIP protocol. The enriched DNA was quantified by real-time PCR using indicated primers. The amount of immunoprecipitated DNA in each sample is represented as signal relative to the total amount of input chromatin, which is equivalent to one.




Fig5: Dot blot analysis of Histone H3 (acetyl K4) on different proteins with Rabbit anti-Histone H3 (acetyl K4) antibody (HA722035) at 1/1,000 dilution. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution for 1 hour at room temperature.

- Lane 1: Unmodified Histone H3 (negative)
- Lane 2: Mono-Methyl-Histone H3 (Lys4) (negative)
- Lane 3: Di-Methyl-Histone H3 (Lys4) (negative)
- Lane 4: Tri-Methyl-Histone H3 (Lys4) (negative)
- Lane 5: Acetyl-Histone H3 (Lys4) (positive)
- Lane 6: Acetyl-Histone H3 (Lys9) (negative)
- Lane 7: Acetyl-Histone H3 (Lys27) (negative)

Proteins loading: 100ng, 25ng, 5ng;

Blocking and dilution buffer: 5% NFD/MTBST;

Exposure time: 50 seconds.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Wani S et al. Human SCP4 is a chromatin-associated CTD phosphatase and exhibits the dynamic translocation during erythroid differentiation. *J Biochem* 160:111-20 (2016).
2. Ni JZ et al. A transgenerational role of the germline nuclear RNAi pathway in repressing heat stress-induced transcriptional activation in *C. elegans*. *Epigenetics Chromatin* 9:3 (2016).

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
H U A B I O
www.huabio.cn