

Anti-GFAP Antibody

R1308-9



Product Type:	Rabbit polyclonal IgG, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IF-Cell, IHC-P, FC
Molecular Wt:	Predicted band size: 50 kDa

Description: GFAP, a class-III intermediate filament, is a cell-specific marker that, during the development of the central nervous system, distinguishes astrocytes from other glial cells. In particular, vimentin filaments are present at early developmental stages, while GFAP filaments are characteristic of differentiated and mature brain astrocytes. In addition, GFAP intermediate filaments are also present in nonmyelin-forming Schwann cells in the peripheral nervous system.

Immunogen: Synthetic peptide within N-terminal human GFAP.

Positive control: Human brain tissue lysates, mouse brain tissue lysates, A172, human brain tissue, rat hippocampus tissue.

Subcellular location: Cytoplasm, Intermediate filament.

Database links: SwissProt: P14136 Human

Recommended Dilutions:

WB	1:2,000
IF-Cell	1:50-1:200
IHC-P	1:200
FC	1:50-1:100

Storage Buffer: 1*PBS (pH7.4), 0.2% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

Purity: Immunogen affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images




Fig1: Western blot analysis on human brain tissue lysates using anti-GFAP rabbit polyclonal antibodies.




Fig2: Western blot analysis of GFAP on mouse brain tissue lysates with Rabbit anti-GFAP antibody (R1308-9) at 1/500 dilution.

Lysates/proteins at 20 µg/Lane.

Predicted band size: 50 kDa

Observed band size: 50 kDa

Exposure time: 1 minute;

10% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody (R1308-9) at 1/500 dilution was used in 5% NFDN/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1:200,000 dilution was used for 1 hour at room temperature.




Fig3: Immunofluorescent staining of A172 cells using anti- GFAP rabbit polyclonal antibody.




Fig4: Immunohistochemical analysis of paraffin-embedded human brain tissue using anti- GFAP rabbit polyclonal antibody.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn



Fig5: Immunohistochemical analysis of paraffin-embedded rat hippocampus tissue with Rabbit anti-GFAP antibody (R1308-9) at 1/800 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (R1308-9) at 1/800 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. "A new splice variant of glial fibrillary acidic protein GFAPepsilon, interacts with the presenilin proteins." Nielsen A.L., Holm I.E., Johansen M., Bonven B., Jorgensen P., Jorgensen A.L. J. Biol. Chem. 277:29983-29991(2002)
2. "Mutations in GFAP, encoding glial fibrillary acidic protein, are associated with Alexander disease." Brenner M., Johnson A.B., Boespflug-Tanguy O., Rodriguez D., Goldman J.E., Messing A. Nat. Genet. 27:117-120(2001)

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn