

Anti-PLGF Antibody [3G3]

EM1701-88



Product Type:	Mouse monoclonal IgG1, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IHC-P, IF-Cell
Molecular Wt:	Predicted band size: 25 kDa
Clone number:	3G3

Description: The onset of angiogenesis is believed to be an early event in tumorigenesis and may facilitate tumor progression and metastasis. Several growth factors with angiogenic activity have been described. These include fibroblast growth factor (FGF), platelet derived growth factor (PDGF), vascular endothelial growth factor (VEGF) and placenta growth factor (PIGF). Like VEGF, several PIGF variants have been shown to arise from alternative mRNA splicings. Evidence has suggested VEGF to be an obligatory component in PIGF signaling. While VEGF homodimers and VEGF/PIGF heterodimers function as potent mediators of mitogenic and chemotactic responses in endothelial cells, PIGF homodimers are effectual only at extremely high concentrations. Indeed, many of the physiological effects attributed to VEGF may actually be a result of VEGF/PIGF. VEGF and PIGF share a common receptor, Flt-1, and may also activate Flk-1/KDR.

Immunogen: Recombinant protein within human PLGF aa 1-221.

Positive control: JAR, LOVO, rat kidney tissue, human placenta tissue, mouse testis tissue.

Subcellular location: Secreted.

Database links: SwissProt: P49763 Human | P49764 Mouse | Q63434 Rat

Recommended Dilutions:

WB	1:500
IF-Cell	1:50
IHC-P	1:50-1:200

Storage Buffer: 1*PBS (pH7.4), 0.2% BSA, 50% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Shipped at 4°C. Store at +4°C short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20°C long term.

Purity: Immunogen affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images




Fig1: Western blot analysis of PLGF on JAR using anti-PLGF antibody at 1/500 dilution.




Fig2: ICC staining PLGF (green) in LOVO cells. The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Fig3: Immunohistochemical analysis of paraffin-embedded rat kidney tissue using anti-PLGF antibody. Counter stained with hematoxylin.




Fig4: Immunohistochemical analysis of paraffin-embedded human placenta tissue using anti-PLGF antibody. Counter stained with hematoxylin.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn




Fig5: Immunohistochemical analysis of paraffin-embedded mouse testis tissue using anti-PLGF antibody. Counter stained with hematoxylin.

Note: All products are “FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE”.

Background References

1. Rolny C et al. HRG inhibits tumor growth and metastasis by inducing macrophage polarization and vessel normalization through downregulation of PlGF. *Cancer Cell* 19:31-44 (2011).
2. Park J E et al. Placenta growth factor. Potentiation of vascular endothelial growth factor bioactivity, in vitro and in vivo, and high affinity binding to Flt-1 but not to Flk-1/KDR. *J Biol Chem* 269:25646-25654 (1994).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn