

# Anti-NM23 Antibody [13C2]

## EM1901-18



<b>Product Type:</b>	Mouse monoclonal IgG1, primary antibodies
<b>Species reactivity:</b>	Human, Mouse, Rat
<b>Applications:</b>	WB, IHC-P, IF-Cell
<b>Molecular Wt:</b>	Predicted band size: 17 kDa
<b>Clone number:</b>	13C2

**Description:** Nucleoside diphosphate kinase A is an enzyme that in humans is encoded by the NME1 gene. It is thought to be a metastasis suppressor. This gene (NME1) was identified because of its reduced mRNA transcript levels in highly metastatic cells. Nucleoside diphosphate kinase (NDK) exists as a hexamer composed of 'A' (encoded by this gene) and 'B' (encoded by NME2) isoforms. Mutations in this gene have been identified in aggressive neuroblastomas. Two transcript variants encoding different isoforms have been found for this gene. Co-transcription of this gene and the neighboring downstream gene (NME2) generates naturally occurring transcripts (NME1-NME2), which encodes a fusion protein consisting of sequence sharing identity with each individual gene product.

**Immunogen:** Synthetic peptide within Human NM23 aa 10-30.

**Positive control:** HEK-293 cell lysate, MCF7 cell lysate, Jurkat cell lysate, Raji cell lysate, HeLa cell lysate, K-562 cell lysate, A549 cell lysate, A431 cell lysate, HepG2 cell lysate, NIH/3T3 cell lysate, C6 cell lysate, mouse kidney tissue lysate, mouse liver tissue lysate, rat brain tissue lysate, mouse kidney tissue, HeLa, NIH/3T3, C6.

**Subcellular location:** Cytoplasm, Nucleus.

**Database links:** SwissProt: P15531 Human | P15532 Mouse | Q05982 Rat

### Recommended Dilutions:

<b>WB</b>	1:2,000-1:5,000
<b>IHC-P</b>	1:2,000
<b>IF-Cell</b>	1:100

**Storage Buffer:** 1\*PBS (pH7.4), 0.2% BSA, 50% Glycerol. Preservative: 0.05% Sodium Azide.

**Storage Instruction:** Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

**Purity:** Immunogen affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

## Images

**Fig1:** Western blot analysis of NM23 on different lysates with Mouse anti-NM23 antibody (EM1901-18) at 1/2,000 dilution.



- Lane 1: HEK-293 cell lysate (20 µg/Lane)
- Lane 2: MCF7 cell lysate (20 µg/Lane)
- Lane 3: Jurkat cell lysate (20 µg/Lane)
- Lane 4: Raji cell lysate (20 µg/Lane)
- Lane 5: HeLa cell lysate (20 µg/Lane)
- Lane 6: K-562 cell lysate (20 µg/Lane)
- Lane 7: A549 cell lysate (20 µg/Lane)
- Lane 8: A431 cell lysate (20 µg/Lane)
- Lane 9: HepG2 cell lysate (20 µg/Lane)
- Lane 10: NIH/3T3 cell lysate (20 µg/Lane)
- Lane 11: C6 cell lysate (20 µg/Lane)
- Lane 12: Mouse kidney tissue lysate (40 µg/Lane)
- Lane 13: Mouse liver tissue lysate (40 µg/Lane)
- Lane 14: Rat brain tissue lysate (40 µg/Lane)


Predicted band size: 17 kDa

Observed band size: 17/20 kDa

Exposure time: 1 minute 59 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody (EM1901-18) at 1/2,000 dilution was used in 5% NFDN/TBST at 4°C overnight. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/50,000 dilution was used for 1 hour at room temperature.



**Fig2:** Immunohistochemical analysis of paraffin-embedded mouse kidney tissue with Mouse anti-NM23 antibody (EM1901-18) at 1/2,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (EM1901-18) at 1/2,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn

**Fig3:** Western blot analysis of NM23 on different lysates with Mouse anti-NM23 antibody (EM1901-18) at 1/2,000 dilution.

Lane 1: A549-si NT cell lysate  
Lane 2: A549-si NM23 cell lysate



Lysates/proteins at 10 µg/Lane.


Predicted band size: 17 kDa  
Observed band size: 17/20 kDa

Exposure time: 17 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (EM1901-18) at 1/2,000 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/50,000 dilution was used for 1 hour at room temperature.

**Fig4:** Immunocytochemistry analysis of HeLa cells labeling NM23 with Mouse anti-NM23 antibody (EM1901-18) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Mouse anti-NM23 antibody (EM1901-18) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 488, HA1125) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

beta Tubulin (ET1602-4, red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 594, HA1122) were used as the secondary antibody at 1/1,000 dilution.

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn


**Fig5:** Immunocytochemistry analysis of NIH/3T3 cells labeling NM23 with Mouse anti-NM23 antibody (EM1901-18) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Mouse anti-NM23 antibody (EM1901-18) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 488, HA1125) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

beta Tubulin (ET1602-4, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Rabbit IgG H&L (iFluor™ 594, HA1122) were used as the secondary antibody at 1/1,000 dilution.

**Fig6:** Immunocytochemistry analysis of C6 cells labeling NM23 with Mouse anti-NM23 antibody (EM1901-18) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Mouse anti-NM23 antibody (EM1901-18) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 488, HA1125) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

beta Tubulin (ET1602-4, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Rabbit IgG H&L (iFluor™ 594, HA1122) were used as the secondary antibody at 1/1,000 dilution.

---

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

---

### Background References

1. Chowdhury D. et al. The exonuclease TREX1 is in the SET complex and acts in concert with NM23-H1 to degrade DNA during granzyme A-mediated cell death. *Mol. Cell* 23:133-142(2006).
2. Fan Z. et al. Tumor suppressor NM23-H1 is a granzyme A-activated DNase during CTL-mediated apoptosis, and the nucleosome assembly protein SET is its inhibitor. *Cell* 112:659-672(2003).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUAABIO  
www.huabio.cn