

Anti-S tag Antibody [G10-C8]

EM50105



Product Type:	Mouse monoclonal IgG1, primary antibodies
Species reactivity:	Species independent
Applications:	WB, ELISA, IP, IF-Cell
Clone number:	G10-C8

Description: S-tag is the name of an oligopeptide derived from pancreatic ribonuclease A (RNase A). If RNase A is digested with subtilisin, a single peptide bond is cleaved, but the resulting two products remain weakly bound to each other and the protein, called ribonuclease S, remains active although each of the two products alone shows no enzymatic activity. The N-terminus of the original RNase A, also called S-peptide, consists of 20 amino acid residues, of which only the first 15 are required for ribonuclease activity. This 15 amino acids long peptide is called S15 or S-tag. The amino acid sequence of the S-tag is: Lys-Glu-Thr-Ala-Ala-Ala-Lys-Phe-Glu-Arg-Gln-His-Met-Asp-Ser. It is believed that the peptide with its abundance of charged and polar residues could improve solubility of proteins it is attached to. Moreover, the peptide alone is thought not to fold into a distinct structure. On DNA-level the S-tag can be attached to the N- or C-terminus of any protein. After gene expression, such a tagged protein can be detected by commercially available antibodies.

Immunogen: Synthetic peptide.

Positive control: S-tagged recombinant protein.

Recommended Dilutions:

WB	1:5,000
ELISA	Use at an assay dependent concentration.
IF-Cell	Use at an assay dependent concentration.
IP	Use at an assay dependent concentration.

Storage Buffer: 1*PBS (pH7.4), 0.2% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Store at +4℃ after thawing. Aliquot store at -20℃ or -80℃. Avoid repeated freeze / thaw cycles.

Purity: Protein G affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images




Fig1: Western blot analysis of S tag on S tag recombinant protein with Mouse anti-S tag antibody (EM50105) at 1/1,000 dilution.

Lysates/proteins at 50 ng/Lane.

Exposure time: 25 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (EM50105) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/50,000 dilution was used for 1 hour at room temperature.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Ning YJ, et al. Viral suppression of innate immunity via spatial isolation of TBK1/IKKe from mitochondrial antiviral platform. *J Mol Cell Biol.* 6:324-37 (2014).
2. Williams JM, et al. The ERdj5-Sel1L complex facilitates cholera toxin retrotranslocation. *Mol Biol Cell.* 24:785-95 (2013).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn