

Anti-CD276 Antibody

ER1803-36



Product Type:	Rabbit polyclonal IgG, primary antibodies
Species reactivity:	Human
Applications:	WB, IF-Cell
Molecular Wt:	Predicted band size: 57 kDa

Description: May participate in the regulation of T-cell-mediated immune response. May play a protective role in tumor cells by inhibiting natural-killer mediated cell lysis as well as a role of marker for detection of neuroblastoma cells. May be involved in the development of acute and chronic transplant rejection and in the regulation of lymphocytic activity at mucosal surfaces. Could also play a key role in providing the placenta and fetus with a suitable immunological environment throughout pregnancy. Both isoform 1 and isoform 2 appear to be redundant in their ability to modulate CD4 T-cell responses. Isoform 2 is shown to enhance the induction of cytotoxic T-cells and selectively stimulates interferon gamma production in the presence of T-cell receptor signaling.

Immunogen: Recombinant protein within human CD276 aa 1-466 / 534.

Positive control: MCF7 cell lysate, HEK-293 cell lysate, LoVo cell lysate, U-2 OS cell lysate, LNCaP cell lysate, SH-SY5Y cell lysate, THP-1 cell lysate, HCT 116 cell lysate, MCF7.

Subcellular location: Membrane.

Database links: SwissProt: Q5ZPR3 Human

Recommended Dilutions:

WB	1:2,000-1:5,000
IF-Cell	1:500

Storage Buffer: PBS (pH7.4), 0.1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Immunogen affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images

Fig1: Western blot analysis of CD276 on different lysates with Rabbit anti-CD276 antibody (ER1803-36) at 1/2,000 dilution.



Lane 1: MCF7 cell lysate
 Lane 2: HEK-293 cell lysate
 Lane 3: LoVo cell lysate
 Lane 4: U-2 OS cell lysate
 Lane 5: LNCaP cell lysate
 Lane 6: SH-SY5Y cell lysate
 Lane 7: THP-1 cell lysate
 Lane 8: HCT 116 cell lysate


Lysates/proteins at 20 µg/Lane.

Predicted band size: 57 kDa
 Observed band size: 100 kDa

Exposure time: 42 seconds; ECL: K1802;
 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (ER1803-36) at 1/2,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig2: Western blot analysis of CD276 on different lysates with Rabbit anti-CD276 antibody (ER1803-36) at 1/5,000 dilution.



Lane 1: HAP1-parental cell lysate
 Lane 2: HAP1-CD276 KD cell lysate

Lysates/proteins at 10 µg/Lane.

Predicted band size: 57 kDa
 Observed band size: 100 kDa

Exposure time: 8 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (ER1803-36) at 1/5,000 dilution was used in K1803 at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
 HUABIO
 www.huabio.cn

Fig3: Immunocytochemistry analysis of MCF7 cells labeling CD276 with Rabbit anti-CD276 antibody (ER1803-36) at 1/500 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-CD276 antibody (ER1803-36) at 1/500 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

Note: All products are “FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE”.

Background References

1. Chapoval A.I et al. B7-H3: a costimulatory molecule for T cell activation and IFN-gamma production. *Nat Immunol* 2:269-274 (2001).
2. Steinberger P et al. Molecular characterization of human 4Ig-B7-H3, a member of the B7 family with four Ig-like domains. *J Immunol* 172:2352-2359 (2004).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn