Anti-RANK Antibody

ER1915-70



Product Type: Rabbit polyclonal IgG, primary antibodies

Species reactivity: Human, Mouse, Rat
Applications: WB, IHC-P, IF-Cell

Molecular Wt: 66 kDa

Description: The protein encoded by this gene is a member of the TNF-receptor superfamily. This

receptors can interact with various TRAF family proteins, through which this receptor induces the activation of NF-kappa B and MAPK8/JNK. This receptor and its ligand are important regulators of the interaction between T cells and dendritic cells. This receptor is

also an essential

Immunogen: Synthetic peptide from human protein at AA range: 60-120

Subcellular location: Membrane.

Database links: SwissProt: Q9Y6Q6 Human

Recommended Dilutions:

WB 1:500-2000 IHC-P 1:200-500 IF-cell 1:50-200

Storage Buffer: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Storage Instruction: -15° to -25° /1 year(Do not lower than -25°)

Purity: Immunogen affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.





Images




Fig1: Western blot analysis of 293T mouse-kidney, antibody was diluted at 1,000. Secondary antibody(catalog#:RS0002) was diluted at 1:20,000




Fig2: Immunohistochemical analysis of paraffin-embedded humanliver, antibody was diluted at 1:200

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".