

Anti-ATP citrate lyase Antibody [ST51-07]

ET1609-37



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IF-Cell, IF-Tissue, IHC-P, FC, IP
Molecular Wt:	Predicted band size: 121 kDa
Clone number:	ST51-07

Description: ATP-citrate synthase, also designated ATP-citrate lyase or citrate cleavage enzyme, is a cytoplasmic homotetramer belonging to the succinate/malate CoA ligase family. The gene coding for this protein maps against chromosome 17q12-q21. ATP-citrate synthase catalyses the formation of acetyl-CoA and oxaloacetate from citrate and CoA. This product, Acetyl-CoA, is necessary for both fatty acid and cholesterol biosynthesis. ATP citrate-lyase is important in the biosynthesis of acetylcholine in nervous tissue.

Immunogen: Synthetic peptide within C-terminal human ATP citrate lyase.

Positive control: HeLa cell lysate, A549 cell lysate, NIH/3T3 cell lysate, Mouse liver tissue lysate, Rat liver tissue lysate, C6 cell lysate, HeLa, NIH/3T3, human kidney tissue, mouse kidney tissue, rat kidney tissue, human thyroid tissue.

Subcellular location: Cytoplasm, cytosol.

Database links: SwissProt: P53396 Human | Q91V92 Mouse | P16638 Rat

Recommended Dilutions:

WB	1:1,000-1:5,000
IF-Cell	1:1,000
IF-Tissue	1:50-1:200
IHC-P	1:200-1:1,000
FC	1:1,000
IP	1-2µg/sample

Storage Buffer: 1*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Shipped at 4°C. Store at +4°C short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20°C long term.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images





Fig1: Western blot analysis of ATP citrate lyase with anti-ATP citrate lyase antibody [ST51-07] (ET1609-37) at 1/1,000 dilution. Lane 1: Wild-type A549 whole cell lysate (20 μ g). Lane 2: ATP citrate lyase knockout A549 whole cell lysate (20 μ g).

ET1609-37 was shown to specifically react with ATP citrate lyase in wild-type A549 cells. No band was observed when ATP citrate lyase knockout sample was tested. Wild-type and ATP citrate lyase knockout samples were subjected to SDS-PAGE. Proteins were transferred to a PVDF membrane and blocked with 5% NFDM in TBST for 1 hour at room temperature. The primary antibody (ET1609-37, 1/1,000) and Loading control antibody (Rabbit anti-HSP90, ET1605-56, 1/10,000) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Rabbit IgG-HRP Secondary Antibody (HA1001) at 1:200,000 dilution was used for 1 hour at room temperature.

Fig2: Western blot analysis of ATP citrate lyase on different lysates with Rabbit anti-ATP citrate lyase antibody (ET1609-37) at 1/5,000 dilution.



Lane 1: HeLa cell lysate (20 μ g/Lane)
 Lane 2: A549 cell lysate (20 μ g/Lane)
 Lane 3: NIH/3T3 cell lysate (20 μ g/Lane)
 Lane 4: Mouse liver tissue lysate (40 μ g/Lane)
 Lane 5: Rat liver tissue lysate (40 μ g/Lane)

Predicted band size: 121 kDa
 Observed band size: 121 kDa

Exposure time: 10 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (ET1609-37) at 1/5,000 dilution was used in primary antibody dilution (K1803) at 4 $^{\circ}$ C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
 HUABIO
 www.huabio.cn

Fig3: Western blot analysis of ATP citrate lyase on different lysates with Rabbit anti-ATP citrate lyase antibody (ET1609-37) at 1/1,000 dilution.

Lane 1: HeLa cell lysate
Lane 2: NIH/3T3 cell lysate
Lane 3: C6 cell lysate



Lysates/proteins at 10 µg/Lane.


Predicted band size: 121 kDa
Observed band size: 121 kDa

Exposure time: 10 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (ET1609-37) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig4: Immunocytochemistry analysis of HeLa cells labeling ATP citrate lyase with Rabbit anti-ATP citrate lyase antibody (ET1609-37) at 1/1,000 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-ATP citrate lyase antibody (ET1609-37) at 1/1,000 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn


Fig5: Immunocytochemistry analysis of NIH/3T3 cells labeling ATP citrate lyase with Rabbit anti-ATP citrate lyase antibody (ET1609-37) at 1/1,000 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-ATP citrate lyase antibody (ET1609-37) at 1/1,000 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.


Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

Fig6: Immunohistochemical analysis of paraffin-embedded human kidney tissue with Rabbit anti-ATP citrate lyase antibody (ET1609-37) at 1/1,000 dilution.



The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ET1609-37) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Fig7: Immunohistochemical analysis of paraffin-embedded mouse kidney tissue with Rabbit anti-ATP citrate lyase antibody (ET1609-37) at 1/1,000 dilution.



The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ET1609-37) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig8: Immunohistochemical analysis of paraffin-embedded rat kidney tissue with Rabbit anti-ATP citrate lyase antibody (ET1609-37) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ET1609-37) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig9: Immunohistochemical analysis of paraffin-embedded human thyroid tissue using anti-ATP citrate lyase antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (ET1609-37, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig10: Flow cytometric analysis of HeLa cells labeling ATP citrate lyase.

Cells were fixed and permeabilized. Then stained with the primary antibody (ET1609-37, 1/1,000) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4°C for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1121) at 1/1,000 dilution for 30 minutes at +4°C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).




Fig11: Flow cytometric analysis of NIH/3T3 cells labeling ATP citrate lyase.

Cells were fixed and permeabilized. Then stained with the primary antibody (ET1609-37, 1/1,000) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4°C for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1121) at 1/1,000 dilution for 30 minutes at +4°C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn




Fig12: ATP citrate lyase was immunoprecipitated from 0.2 mg NIH/3T3 cell lysate with ET1609-37 at 2 μ g/10 μ l beads. Western blot was performed from the immunoprecipitate using ET1609-37 at 1/1,000 dilution. HRP Conjugated Anti-Rabbit IgG for IP Nano-secondary antibody at 1/5,000 dilution was used for 1 hour at room temperature.

Lane 1: NIH/3T3 cell lysate (input)

Lane 2: ET1609-37 IP in NIH/3T3 cell lysate

Lane 3: Rabbit IgG instead of ET1609-37 in NIH/3T3 cell lysate

Blocking/Dilution buffer: primary antibody dilution (K1803)

Exposure time: 2 seconds; ECL: K1801

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

- Walsh J et al. Identification and quantification of the basal and inducible Nrf2-dependent proteomes in mouse liver: biochemical, pharmacological and toxicological implications. *J Proteomics* 108:171-87 (2014).
- Dani Is VW et al. Cancer cells differentially activate and thrive on de novo lipid synthesis pathways in a low-lipid environment. *PLoS One* 9:e106913 (2014).

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn