

# Anti-BCL2A1 Antibody [SC05-42]

ET1610-20



<b>Product Type:</b>	Recombinant Rabbit monoclonal IgG, primary antibodies
<b>Species reactivity:</b>	Human, Mouse
<b>Applications:</b>	WB, IHC-P
<b>Molecular Wt:</b>	Predicted band size: 20 kDa
<b>Clone number:</b>	SC05-42

**Description:** The Bcl-2 family of proteins is characterized by its ability to modulate cell death under a broad range of physiological conditions. Bcl-2 and Bcl-xL function to inhibit apoptosis while other members of the Bcl-2 family, Bax, Bad, Bak and Bcl-xS, oppose death-suppressing effects. An additional member of the family, A1 (also designated Bfl-1), dimerizes with both Bcl-2 and Bax and has been identified as a hematopoietic- specific, early inducible gene. While A1 demonstrates life promoting properties similar to those of Bcl-2, its function may be more temporally regulated during myeloid differentiation and dependent on additional growth stimuli to confer its life promoting properties. A1 is abundantly expressed in bone marrow and at low levels in other tissues. There is evidence that a correlation exists between a high expression of the A1 gene product and stomach cancer.

**Immunogen:** Synthetic peptide within Human BCL2A1 aa 2-48 / 175.

**Positive control:** Human tonsil tissue, human kidney tissue, mouse liver tissue, mouse kidney tissue, 293, human liver tissue, human kidney tissue lysates.

**Subcellular location:** Cytoplasm.

**Database links:** SwissProt: Q16548 Human | Q07440 Mouse  
Unigene: 19770 Rat

**Recommended Dilutions:**

<b>WB</b>	1:1,000
<b>IHC-P</b>	1:50-1:400

**Storage Buffer:** 1\*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

**Storage Instruction:** Shipped at 4°C. Store at +4°C short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20°C long term.

**Purity:** Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

## Images



**Fig1:** Western blot analysis of BCL2A1 on human kidney tissue lysates with Rabbit anti-BCL2A1 antibody (ET1610-20) at 1/500 dilution.

Lysates/proteins at 40 µg/Lane.


Predicted band size: 20 kDa

Observed band size: 26 kDa

Exposure time: 40 seconds;


4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFD/MTBST for 1 hour at room temperature. The primary antibody (ET1610-20) at 1/500 dilution was used in 5% NFD/MTBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1:100,000 dilution was used for 1 hour at room temperature.



**Fig2:** Immunohistochemical analysis of paraffin-embedded human tonsil tissue with Rabbit anti-BCL2A1 antibody (ET1610-20) at 1/50 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1610-20) at 1/50 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig3:** Immunohistochemical analysis of paraffin-embedded human kidney tissue with Rabbit anti-BCL2A1 antibody (ET1610-20) at 1/400 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1610-20) at 1/400 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn




**Fig4:** Immunohistochemical analysis of paraffin-embedded mouse liver tissue with Rabbit anti-BCL2A1 antibody (ET1610-20) at 1/100 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1610-20) at 1/100 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig5:** Immunohistochemical analysis of paraffin-embedded mouse kidney tissue with Rabbit anti-BCL2A1 antibody (ET1610-20) at 1/400 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1610-20) at 1/400 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig6:** Immunohistochemical analysis of paraffin-embedded human liver tissue with Rabbit anti-BCL2A1 antibody (ET1610-20) at 1/400 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (ET1610-20) at 1/400 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

---

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

---

### Background References

1. Jauharoh SN et al. SS-A/Ro52 promotes apoptosis by regulating Bcl-2 production. *Biochem Biophys Res Commun* 417:582-7 (2012).
2. Fardin P et al. Induction of epithelial mesenchymal transition and vasculogenesis in the lenses of Dbl oncogene transgenic mice. *PLoS One* 4:e7058 (2009).

**Hangzhou Huan Biotechnology Co., Ltd.**

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
H U A B I O  
www.huabio.cn