

# HRP Conjugated Goat Anti-human IgG polyclonal Antibody

## HA1018



<b>Product Type:</b>	Goat polyclonal IgG, secondary antibodies
<b>Species reactivity:</b>	Human
<b>Applications:</b>	WB, ELISA

**Description:** Whole IgG antibodies are isolated as intact molecules from antisera by immunoaffinity chromatography. They have an Fc portion and two antigen binding Fab portions joined together by disulfide bonds and therefore they are divalent. The average molecular weight is reported to be about 160 kDa. The whole IgG form of antibodies is suitable for the majority of immunodetection procedures and is the most cost effective. Horseradish peroxidase (HRP) conjugates are prepared by a modified Nakane and Kawaoi procedure (J. Histochem. Cytochem. 1974. 22, 1084). Peroxidase conjugates are commonly used for immunohistochemistry, Western blotting, and ELISA. Affinity-purified anti-horseradish peroxidase and conjugates are available for detection of horseradish peroxidase antigen or for signal amplification of HRP-containing reagents. For immunostaining of mammalian cells, an advantage of using anti-horseradish peroxidase is reduced background, since the antibody does not recognize the endogenous peroxidase-like enzymes found in those cells.

**Conjugate:** HRP-conjugated

**Immunogen:** Human IgG.

**Positive control:** Human serum lysates.

**Recommended Dilutions:**

<b>WB</b>	1:1000-1:5000
<b>ELISA</b>	1:2000-1:5000

**Storage Buffer:** 1\*TBS (pH7.4), 1% BSA, 50% Glycerol.

**Storage Instruction:** Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

**Purity:** Human IgG affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

## Images



**Fig1:** Western blot analysis of Rabbit anti Goat Anti-human IgG-HRP on Human serum lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. Rabbit anti Goat IgG-HRP Secondary Antibody (HA1018) at 1:2,000 (Lane 1), 1:5,000 (Lane 2) dilution was used for 2 hour at room temperature.

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Hangzhou Huaan Biotechnology Co., Ltd.

Orders: 0086-571-88062880

Technical: 0086-571-89986345

Service mail: support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn