

# iFluor™ 594 Conjugated Goat anti-rabbit IgG polyclonal Antibody

## HA1122



<b>Product Type:</b>	Goat polyclonal IgG, secondary antibodies
<b>Species reactivity:</b>	Rabbit
<b>Applications:</b>	IF-Cell, IF-Tissue, FC, IHC-Fr

**Description:** Whole IgG antibodies are isolated as intact molecules from antisera by immunoaffinity chromatography. They have an Fc portion and two antigen binding Fab portions joined together by disulfide bonds and therefore they are divalent. The average molecular weight is reported to be about 160 kDa. The whole IgG form of antibodies is suitable for the majority of immunodetection procedures and is the most cost effective. iFluor™ 594 dyes have fluorescence excitation and emission maxima of ~590 nm and ~610 nm respectively. iFluor™ 594 family is pH-independent from pH 3 to 11. These spectral characteristics make this new dye family an excellent alternative. iFluor™ 594 is much easier to be conjugated with RPE with much higher conjugation yield, and the resulted RPE-iFluor™ 594 tandem has better FRET efficiency. iFluor™ 594 SE is reasonably stable and shows good reactivity and selectivity with protein amino groups.

**Conjugate:** iFluor™ 594, Ex: 588nm; Em: 604nm.

**Immunogen:** Rabbit IgG (H+L).

**Recommended Dilutions:**

<b>IF-Cell</b>	1:500-1:1,000
<b>IF-Tissue</b>	1:500-1:1,000
<b>FC</b>	1:500-1:1,000
<b>IHC-Fr</b>	1:500-1:1,000

**Storage Buffer:** 1\*PBS (pH7.4), 1% BSA, 40% Glycerol, 0.2% Proclean 950.

**Storage Instruction:** Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

**Purity:** Immunogen affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn


## Images



**Fig1:** Immunofluorescence analysis of paraffin-embedded mouse brain tissue labeling Synaptophysin (ET1606-56) and beta III tubulin (M0805-8).


The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 10% negative goat serum for 1 hour at room temperature, washed with PBS. And then probed with the primary antibodies Synaptophysin (ET1606-56, red) at 1/200 dilution and beta III tubulin (M0805-8, green) at 1/200 dilution at +4°C overnight, washed with PBS. Goat Anti-Rabbit IgG H&L (iFluor™ 594, HA1122) and Goat Anti-Mouse IgG H&L (iFluor™ 488, HA1125) were used as the secondary antibodies at 1/1,000 dilution. DAPI was used as nuclear counterstain.

**Fig2:** Immunocytochemistry analysis of HepG2 cells labeling eIF-6 with Mouse anti-eIF-6 antibody (HA600051) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Mouse anti-eIF-6 antibody (HA600051) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 488, HA1125) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

beta Tubulin (ET1602-4, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Rabbit IgG H&L (iFluor™ 594, HA1122) were used as the secondary antibody at 1/1,000 dilution.



**Fig3:** Flow cytometric analysis of HeLa cells labeling Lamin A/C.

Cells were fixed and permeabilized. Then stained with the primary antibody (ET7110-12, 1ug/ml) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4°C for an hour, the cells were stained with a iFluor™ 594 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1122) at 1/1,000 dilution for 30 minutes at +4°C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).


Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn



**Fig4:** Flow cytometric analysis of K-562 cells labeling LFNG.

Cells were fixed and permeabilized. Then stained with the primary antibody (HA721367, 1ug/ml) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4 °C for an hour, the cells were stained with a iFluor™ 594 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1122) at 1/1,000 dilution for 30 minutes at +4 °C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn