

iFluor™ 750 Conjugated Goat anti-Chicken IgY Antibody

HA1145



Product Type:	Goat polyclonal IgG, secondary antibodies
Species reactivity:	Chicken
Applications:	FC, IHC-Fr, IF-Tissue, IF-Cell

Description: Whole IgG antibodies are isolated as intact molecules from antisera by immunoaffinity chromatography. They have an Fc portion and two antigen binding Fab portions joined together by disulfide bonds and therefore they are divalent. The average molecular weight is reported to be about 160 kDa. The whole IgG form of antibodies is suitable for the majority of immunodetection procedures and is the most cost effective.

Conjugate: iFluor™ 750

Immunogen: Recombinant protein.

Recommended Dilutions:

FC	1:1,000
IHC-Fr	1:500
IF-Tissue	1:500
IF-Cell	1:500

Storage Buffer: Preservative: 0.02% Sodium azide Constituents: 30% Glycerol, 1% BSA, 68.98% PBS

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Immunogen affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images




Fig1: Flow cytometric analysis of THP-1 cells labeling Iba1.

Cells were fixed and permeabilized. Then stained with the primary antibody (HA601376, 1/1,000) (red). After incubation of the primary antibody at +4°C for an hour, the cells were stained with a iFluor™ 750 conjugate-Goat anti-Chicken IgY Secondary antibody (HA1145) at 1/1,000 dilution for 30 minutes at +4°C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).

Note: All products are “FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE”.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn