

Human CD30 Ligand / TNFSF8, Tag Free Protein

HA211293



Product name:	Human CD30 Ligand / TNFSF8, Tag Free
Species reactivity:	Human
Bio-Activity:	Testing in progress.
Protein construction description:	A DNA sequence encoding the human CD30 Ligand / TNFSF8 protein (P32971) (Gln 63-Asp 234) was expressed with tag free.

Background: CD153 (cluster of differentiation 153) also known as tumor necrosis factor ligand superfamily member 8 is a protein that in humans is encoded by the TNFSF8 gene. CD153 is a cytokine ligand for the TNF receptor CD30. It plays a role in the T cell-dependent anti-mycobacterial immune response.

Purity: >95% as determined by SDS-PAGE.

Endotoxin: Less than 1.0 EU per µg by the LAL method.

Fragment region: CD30 Ligand / TNFSF8 (63-234)

Source: HEK293

Accession: P32971

Predicted molecular mass: 20.4 kD

Formulation: Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4, 5% Trehalose, 5% mannitol.

Reconstitution: Reconstitute at 250 µg/ml in sterile water.

Storage: Please avoid repeated freeze-thaw cycles. Samples are stable for up to twelve months from date of receipt at -20°C to -80°C. It is recommended that aliquot the reconstituted solution to minimize freeze-thaw cycles.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders: 0086-571-88062880

Technical: 0086-571-89986345

Service mail: support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Applications: WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation




Fig1: Protein on SDS-PAGE under reducing (R) condition.

Note: All products are “FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE”.