

Anti-NM23 Antibody [13C1-R]

HA601238



Product Type:	Recombinant Mouse monoclonal IgG1, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IF-Cell
Molecular Wt:	Predicted band size: 17 kDa
Clone number:	13C1-R

Description: Major role in the synthesis of nucleoside triphosphates other than ATP. The ATP gamma phosphate is transferred to the NDP beta phosphate via a ping-pong mechanism, using a phosphorylated active-site intermediate. Possesses nucleoside-diphosphate kinase, serine/threonine-specific protein kinase, geranyl and farnesyl pyrophosphate kinase, histidine protein kinase and 3'-5' exonuclease activities. Involved in cell proliferation, differentiation and development, signal transduction, G protein-coupled receptor endocytosis, and gene expression. Required for neural development including neural patterning and cell fate determination. During GZMA-mediated cell death, works in concert with TREX1. NME1 nicks one strand of DNA and TREX1 removes bases from the free 3' end to enhance DNA damage and prevent DNA end reannealing and rapid repair.

Immunogen: Synthetic peptide within Human NM23 aa 1-50 / 152.

Positive control: HEK-293 cell lysate, MCF7 cell lysate, Jurkat cell lysate, HeLa cell lysate, HepG2 cell lysate, K-562 cell lysate, A549 cell lysate, Raji cell lysate, NIH/3T3 cell lysate, C6 cell lysate, mouse kidney tissue lysate, mouse liver tissue lysate, mouse brain tissue lysate, rat brain tissue lysate, rat liver tissue lysate, MCF7.

Subcellular location: Nucleus, Cytoplasm.

Database links: SwissProt: P15531 Human | P15532 Mouse | Q05982 Rat

Recommended Dilutions:

WB	1:1,000
IF-Cell	1:100

Storage Buffer: PBS (pH7.4), 0.1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images

Fig1: Western blot analysis of NM23 on different lysates with Mouse anti-NM23 antibody (HA601238) at 1/1,000 dilution.



Lane 1: HEK-293 cell lysate (20 µg/Lane)

Lane 2: MCF7 cell lysate (20 µg/Lane)

Lane 3: Jurkat cell lysate (20 µg/Lane)

Lane 4: HeLa cell lysate (20 µg/Lane)

Lane 5: HepG2 cell lysate (20 µg/Lane)

Lane 6: K-562 cell lysate (20 µg/Lane)

Lane 7: A549 cell lysate (20 µg/Lane)

Lane 8: Raji cell lysate (20 µg/Lane)

Predicted band size: 17 kDa


Observed band size: 17/20 kDa

Exposure time: 3 minutes; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDm/TBST for 1 hour at room temperature. The primary antibody (HA601238) at 1/1,000 dilution was used in 5% NFDm/TBST at 4°C overnight. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig2: Western blot analysis of NM23 on different lysates with Mouse anti-NM23 antibody (HA601238) at 1/1,000 dilution.



Lane 1: NIH/3T3 cell lysate (20 µg/Lane)

Lane 2: C6 cell lysate (20 µg/Lane)

Lane 3: Mouse kidney tissue lysate (40 µg/Lane)

Lane 4: Mouse liver tissue lysate (40 µg/Lane)

Lane 5: Mouse brain tissue lysate (40 µg/Lane)

Lane 6: Rat brain tissue lysate (40 µg/Lane)

Lane 7: Rat liver tissue lysate (40 µg/Lane)

Predicted band size: 17 kDa

Observed band size: 17 kDa

Exposure time: 25 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDm/TBST for 1 hour at room temperature. The primary antibody (HA601238) at 1/1,000 dilution was used in 5% NFDm/TBST at 4°C overnight. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/50,000 dilution was used for 1 hour at room temperature.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Fig3: Western blot analysis of NM23 on different lysates with Mouse anti-NM23 antibody (HA601238) at 1/2,000 dilution.

Lane 1: A549-si NT cell lysate
Lane 2: A549-si NM23 cell lysate



Lysates/proteins at 10 µg/Lane.


Predicted band size: 17 kDa
Observed band size: 17/20 kDa

Exposure time: 30 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA601238) at 1/2,000 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig4: Immunocytochemistry analysis of MCF7 cells labeling NM23 with Mouse anti-NM23 antibody (HA601238) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Mouse anti-NM23 antibody (HA601238) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 488, HA1125) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

beta Tubulin (ET1602-4, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Rabbit IgG H&L (iFluor™ 594, HA1122) were used as the secondary antibody at 1/1,000 dilution.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Fan Z. et. al. Tumor suppressor NM23-H1 is a granzyme A-activated DNase during CTL-mediated apoptosis, and the nucleosome assembly protein SET is its inhibitor. *Cell* 112:659-672(2003).
2. Chowdhury D. et. al. The exonuclease TREX1 is in the SET complex and acts in concert with NM23-H1 to degrade DNA during granzyme A-mediated cell death. *Mol. Cell* 23:133-142(2006).

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
H U A B I O
www.huabio.cn