

# Anti-GOT1 Antibody [A2D6-R]

HA601279



<b>Product Type:</b>	Recombinant Mouse monoclonal IgG1, primary antibodies
<b>Species reactivity:</b>	Human, Mouse, Rat
<b>Applications:</b>	WB
<b>Molecular Wt:</b>	Predicted band size: 46 kDa
<b>Clone number:</b>	A2D6-R

**Description:** Biosynthesis of L-glutamate from L-aspartate or L-cysteine. Important regulator of levels of glutamate, the major excitatory neurotransmitter of the vertebrate central nervous system. Acts as a scavenger of glutamate in brain neuroprotection. The aspartate aminotransferase activity is involved in hepatic glucose synthesis during development and in adipocyte glyceroneogenesis. Using L-cysteine as substrate, regulates levels of mercaptopyruvate, an important source of hydrogen sulfide. Mercaptopyruvate is converted into H<sub>2</sub>S via the action of 3-mercaptopyruvate sulfurtransferase (3MST). Hydrogen sulfide is an important synaptic modulator and neuroprotectant in the brain.

**Immunogen:** Recombinant protein within Human GOT1 aa 16-211 / 413.

**Positive control:** HeLa cell lysate, K-562 cell lysate, HL-60 cell lysate, Caco-2 cell lysate, human liver tissue lysate, mouse liver tissue lysate, mouse heart tissue lysate, rat liver tissue lysate, rat heart tissue lysate.

**Subcellular location:** Cytoplasm.

**Database links:** SwissProt: P17174 Human | P05201 Mouse | P13221 Rat

**Recommended Dilutions:**

**WB** 1:1,000

**Storage Buffer:** PBS (pH7.4), 0.1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

**Storage Instruction:** Shipped at 4°C. Store at +4°C short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20°C long term.

**Purity:** Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

## Images

**Fig1:** Western blot analysis of GOT1 on different lysates with Mouse anti-GOT1 antibody (HA601279) at 1/1,000 dilution.



Lane 1: HeLa cell lysate (20 µg/Lane)  
 Lane 2: K-562 cell lysate (20 µg/Lane)  
 Lane 3: HL-60 cell lysate (20 µg/Lane)  
 Lane 4: Caco-2 cell lysate (20 µg/Lane)  
 Lane 5: Human liver tissue lysate (40 µg/Lane)  
 Lane 6: Mouse liver tissue lysate (40 µg/Lane)  
 Lane 7: Mouse heart tissue lysate (40 µg/Lane)  
 Lane 8: Rat liver tissue lysate (40 µg/Lane)  
 Lane 9: Rat heart tissue lysate (40 µg/Lane)

Lysates/proteins at 10 µg/Lane.


Predicted band size: 46 kDa  
 Observed band size: 38 kDa

Exposure time: 1 minute 41 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDm/TBST for 1 hour at room temperature. The primary antibody (HA601279) at 1/1,000 dilution was used in 5% NFDm/TBST at 4°C overnight. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/50,000 dilution was used for 1 hour at room temperature.

**Fig2:** Western blot analysis of GOT1 on different lysates with Mouse anti-GOT1 antibody (HA601279) at 1/1,000 dilution.



Lane 1: HAP1-parental cell lysate  
 Lane 2: HAP1-GOT1 KD cell lysate

Lysates/proteins at 10 µg/Lane.

Predicted band size: 43 kDa  
 Observed band size: 43 kDa

Exposure time: 2 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDm/TBST for 1 hour at room temperature. The primary antibody (HA601279) at 1/1,000 dilution was used in K1803 at 4°C overnight. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/50,000 dilution was used for 1 hour at room

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
 HUABIO  
 www.huabio.cn

---

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

---

### Background References

1. Abrego J. et. al. GOT1-mediated anaplerotic glutamine metabolism regulates chronic acidosis stress in pancreatic cancer cells. *Cancer Lett.* 2017 Aug 1;400:37-46.
2. Hong C. et. al. Inhibition of GOT1 sensitizes colorectal cancer cells to 5-fluorouracil. *Cancer Chemother Pharmacol.* 2017 Apr;79(4):835-840.

**Hangzhou Huaan Biotechnology Co., Ltd.**

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUAABIO  
www.huabio.cn