

# Anti-p53 Antibody [6B5-R]

## HA601315



<b>Product Type:</b>	Recombinant Mouse monoclonal IgG1, primary antibodies
<b>Species reactivity:</b>	Human, Mouse, Monkey
<b>Applications:</b>	WB, IF-Cell
<b>Molecular Wt:</b>	Predicted band size: 53 kDa
<b>Clone number:</b>	6B5-R

**Description:** Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Induces the transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA-Mkl1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and seem to have to effect on cell-cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis. Regulates the circadian clock by repressing CLOCK-ARNTL/BMAL1-mediated transcriptional activation of PER2.

<b>Immunogen:</b>	Recombinant protein within C-terminal human p53.
<b>Positive control:</b>	293T cell lysate, MCF7 cell lysate, NIH/3T3 treated with 0.5 $\mu$ M doxorubicin cell for 24 hours lysate, Neuro-2a cell lysate, COS-1 cell lysate, A431.
<b>Subcellular location:</b>	Nucleus. Cytoplasm. Localized in both nucleus and cytoplasm in most cells. In some cells, forms foci in the nucleus that are different from nucleoli.
<b>Database links:</b>	SwissProt: P04637 Human   P02340 Mouse
<b>Recommended Dilutions:</b>	
<b>WB</b>	1:1,000-1:2,000
<b>IF-Cell</b>	1:100
<b>Storage Buffer:</b>	PBS (pH7.4), 0.1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.
<b>Storage Instruction:</b>	Store at +4 $^{\circ}$ C after thawing. Aliquot store at -20 $^{\circ}$ C. Avoid repeated freeze / thaw cycles.
<b>Purity:</b>	Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

## Images

**Fig1:** Western blot analysis of p53 on different lysates with Mouse anti-p53 antibody (HA601315) at 1/1,000 dilution.

Lane 1: 293T cell lysate  
 Lane 2: MCF7 cell lysate  
 Lane 3: NIH/3T3 cell lysate  
 Lane 4: NIH/3T3 treated with 0.5 $\mu$ M doxorubicin for 24 hours lysate  
 Lane 5: Neuro-2a cell lysate  
 Lane 6: COS-1 cell lysate



Lysates/proteins at 20  $\mu$ g/Lane.

Predicted band size: 53 kDa  
 Observed band size: 53 kDa


Exposure time: Lane 1-2: 1 minute 40 seconds; Lane 3-6: 3 minutes;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA601315) at 1/1,000 dilution was used in 5% NFDM/TBST at 4 $^{\circ}$ C overnight. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/50,000 dilution was used for 1 hour at room temperature.

**Fig2:** Western blot analysis of p53 on different lysates with Mouse anti-p53 antibody (HA601315) at 1/2,000 dilution.

Lane 1: HAP1-parental cell lysate  
 Lane 2: HAP1-p53 KD cell lysate



Lysates/proteins at 10  $\mu$ g/Lane.

Predicted band size: 53 kDa  
 Observed band size: 53 kDa

Exposure time: 40 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA601315) at 1/2,000 dilution was used in K1803 at 4 $^{\circ}$ C overnight. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/50,000 dilution was used for 1 hour at room temperature.

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
 HUABIO  
 www.huabio.cn

**Fig3:** Immunocytochemistry analysis of A431 cells labeling p53 with Mouse anti-p53 antibody (HA601315) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Mouse anti-p53 antibody (HA601315) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 488, HA1125) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

beta Tubulin (ET1602-4, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Rabbit IgG H&L (iFluor™ 594, HA1122) were used as the secondary antibody at 1/1,000 dilution.

**Note:** All products are “FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE”.

### Background References

1. Louria-Hayon I et al. The promyelocytic leukemia protein protects p53 from Mdm2-mediated inhibition and degradation. *J Biol Chem* 278:33134-33141 (2003).
2. An W et al. Ordered cooperative functions of PRMT1, p300, and CARM1 in transcriptional activation by p53. *Cell* 117:735-748 (2004).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn