

TDP43 Recombinant Antibody [PSH09-14] - Rat IgG1 (Chimeric)

HA601627



Product Type:	Recombinant Chimeric Antibody, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IHC-P, IF-Tissue, IF-Cell
Molecular Wt:	Predicted band size: 45 kDa
Clone number:	PSH09-14

Description: TDP-43 is a transcriptional repressor that binds to chromosomally integrated TAR DNA and represses HIV-1 transcription. In addition, this protein regulates alternate splicing of the CFTR gene. TDP-43 has been shown to bind both DNA and RNA and have multiple functions in transcriptional repression, pre-mRNA splicing and translational regulation. Recent work has characterized the transcriptome-wide binding sites revealing that thousands of RNAs are bound by TDP-43 in neurons. TDP-43 was originally identified as a transcriptional repressor that binds to chromosomally integrated trans-activation response element (TAR) DNA and represses HIV-1 transcription. It was also reported to regulate alternate splicing of the CFTR gene and the apoA-II gene. In spinal motor neurons TDP-43 has also been shown in humans to be a low molecular weight neurofilament (hNFL) mRNA-binding protein. It has also shown to be a neuronal activity response factor in the dendrites of hippocampal neurons suggesting possible roles in regulating mRNA stability, transport and local translation in neurons. It has been demonstrated that zinc ions are able to induce aggregation of endogenous TDP-43 in cells. Moreover, zinc could bind to RNA binding domain of TDP-43 and induce the formation of amyloid-like aggregates in vitro.

Immunogen: Synthetic peptide within human TDP43 aa 365-414.

Positive control: HeLa (Human cervical adenocarcinoma cell) lysate, HEK-293 (Human embryonic kidney cell) lysate, SH-SY5Y (Human neuroblastoma cell) lysate, K-562 (Human chronic myelogenous leukemia cell) lysate, C2C12 (Mouse myoblast) lysate, PC-12 (Rat pheochromocytoma cell (undifferentiated)) lysate, Mouse brain tissue lysate.

Subcellular location: Cytoplasm, Mitochondrion, Nucleus.

Database links: SwissProt: Q13148 Human | Q921F2 Mouse
Entrez Gene: 298648 Rat

Recommended Dilutions:

WB	1:2,000
IHC-P	1:500-1:2,000
IF-Tissue	1:500
IF-Cell	1:50

Storage Buffer: 1*PBS (pH7.4), 0.1% BSA, 40% Glycerol, 0.2% Proclean 950.

Storage Instruction: Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C long term.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images

Fig1: Western blot analysis of TDP43 on different lysates with Rat anti-TDP43 antibody (HA601627) at 1/2,000 dilution.



Lane 1: HeLa (Human cervical adenocarcinoma cell) lysate

Lane 2: HEK-293 (Human embryonic kidney cell) lysate

Lane 3: SH-SY5Y (Human neuroblastoma cell) lysate

Lane 4: K-562 (Human chronic myelogenous leukemia cell) lysate

Lane 5: C2C12 (Mouse myoblast) lysate

Lane 6: PC-12 (Rat pheochromocytoma cell (undifferentiated)) lysate

Lane 7: Mouse brain tissue lysate

Lysates/proteins at 15 µg/Lane.

Exposure time: 20 seconds; ECL: K1801

Blocking: 5% NFDm/TBST, 1 hour at room temperature

Primary antibody: HA601627, 1/2,000 in primary antibody dilution buffer (K1803), overnight at 4 °C

Secondary antibody: Goat anti-Rat IgG-HRP (HA1023), 1/50,000 in 5% NFDm/TBST, 1 hour at room temperature

Predicted band size: 45 kDa


Observed band size: 45 kDa

Fig2: Application: Immunohistochemistry (IHC-P)

Species: Human

Tissue: Brain

Sample: Paraffin-embedded section



Antigen retrieval: Heat-mediated, Tris-EDTA buffer (pH 9.0), 20 minutes at 95°C.

Wash buffer: 1× TBST

Endogenous peroxidase blocking: 3% H₂O₂, 10 minutes at room temperature.

Blocking: 1% BSA + 10% normal goat serum, 10 minutes at room temperature.

Primary antibody: HA601627, 1/500, 1 hour at room temperature.

Secondary antibody: HA1023, 30 minutes at room temperature.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn




Fig3: Application: Immunohistochemistry (IHC-P)

Species: Mouse

Tissue: Brain

Sample: Paraffin-embedded section

Antigen retrieval: Heat-mediated, Tris-EDTA buffer (pH 9.0), 20 minutes at 95°C.

Wash buffer: 1× TBST

Endogenous peroxidase blocking: 3% H₂O₂, 10 minutes at room temperature.

Blocking: 1% BSA + 10% normal goat serum, 10 minutes at room temperature.

Primary antibody: HA601627, 1/2,000, 1 hour at room temperature.

Secondary antibody: HA1023, 30 minutes at room temperature.




Fig4: Application: Immunohistochemistry (IHC-P)

Species: Rat

Tissue: Brain

Sample: Paraffin-embedded section

Antigen retrieval: Heat-mediated, Tris-EDTA buffer (pH 9.0), 20 minutes at 95°C.

Wash buffer: 1× TBST

Endogenous peroxidase blocking: 3% H₂O₂, 10 minutes at room temperature.

Blocking: 1% BSA + 10% normal goat serum, 10 minutes at room temperature.

Primary antibody: HA601627, 1/2,000, 1 hour at room temperature.

Secondary antibody: HA1023, 30 minutes at room temperature.




Fig5: Application: Immunofluorescence (IF-tissue)

Species: Mouse

Tissue: Brain

Sample: Paraffin-embedded section

Antigen retrieval: Heat-mediated, Tris-EDTA buffer (pH 9.0), 20 minutes at 95°C.

Wash buffer: 1× TBST

Blocking: 10% normal goat serum + 1% Triton X-100 + 0.3 M Glycine in TBST, 30 minutes at room temperature.

Primary antibody: HA601627, 1/500, overnight at 4°C.

Secondary antibody: Goat Anti-Rat IgG (iFluor™ 488, HA1133), 1.5 hours at room temperature.




Fig6: Application: Immunofluorescence (IF-tissue)

Species: Rat

Tissue: Brain

Sample: Paraffin-embedded section

Antigen retrieval: Heat-mediated, Tris-EDTA buffer (pH 9.0), 20 minutes at 95°C.

Wash buffer: 1× TBST

Blocking: 10% normal goat serum + 1% Triton X-100 + 0.3 M Glycine in TBST, 30 minutes at room temperature.


Primary antibody: HA601627, 1/500, overnight at 4°C.

Secondary antibody: Goat Anti-Rat IgG (iFluor™ 488, HA1133), 1.5 hours at room temperature.

Fig7: Application: Immunocytochemistry (IF-cell)

Species: Human

Sample: HeLa (Human cervix adenocarcinoma epithelial cell)



Fixation: 4% Paraformaldehyde, 15 minutes at room temperature.
 Permeabilization: 0.1% Triton X-100, 15 minutes at room temperature.

Blocking: 1% BSA + 10% normal goat serum, 1 hour at room temperature.

Antibody dilution buffer: 1% BSA in PBST.

Primary antibody: HA601627, 1/50, overnight at 4°C.


Secondary antibody: Goat Anti-Rat IgG (iFluor™ 488, HA1133), 45 minutes at room temperature.

Counterstain: Beta tubulin (ET1602-4, Red), 1/200, overnight at 4°C. The nuclear counterstain was DAPI (Blue).

Fig8: Application: Immunocytochemistry (IF-cell)

Species: Mouse

Sample: C2C12 (Mouse myoblast)



Fixation: 4% Paraformaldehyde, 15 minutes at room temperature.
 Permeabilization: 0.1% Triton X-100, 15 minutes at room temperature.

Blocking: 1% BSA + 10% normal goat serum, 1 hour at room temperature.

Antibody dilution buffer: 1% BSA in PBST.

Primary antibody: HA601627, 1/50, overnight at 4°C.

Secondary antibody: Goat Anti-Rat IgG (iFluor™ 488, HA1133), 45 minutes at room temperature.

Counterstain: Beta tubulin (ET1602-4, Red), 1/200, overnight at 4°C. The nuclear counterstain was DAPI (Blue).

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. McMillan M et al. RNA methylation influences TDP43 binding and disease pathogenesis in models of amyotrophic lateral sclerosis and frontotemporal dementia. *Mol Cell*. 2023 Jan
2. Corbet GA et al. TDP43 ribonucleoprotein granules: physiologic function to pathologic aggregates. *RNA Biol*. 2021 Oct

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn