

Parkin Recombinant Antibody [PSH08-11] - Guinea pig IgG2 (Chimeric)

HA601663



Product Type:	Recombinant Chimeric Antibody, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IF-Cell, IHC-P, IF-Tissue, FC(Intra)
Molecular Wt:	Predicted band size: 52 kDa
Clone number:	PSH08-11

Description: Parkin is a 465-amino acid residue E3 ubiquitin ligase, a protein that in humans and mice is encoded by the PARK2 gene. Parkin plays a critical role in ubiquitination – the process whereby molecules are covalently labelled with ubiquitin (Ub) and directed towards degradation in proteasomes or lysosomes. Ubiquitination involves the sequential action of three enzymes. First, an E1 ubiquitin-activating enzyme binds to inactive Ub in eukaryotic cells via a thioester bond and mobilises it in an ATP-dependent process. Ub is then transferred to an E2 ubiquitin-conjugating enzyme before being conjugated to the target protein via an E3 ubiquitin ligase. There exists a multitude of E3 ligases, which differ in structure and substrate specificity to allow selective targeting of proteins to intracellular degradation.

Immunogen: Recombinant protein within human Parkin aa 1-200.

Positive control: SH-SY5Y (Human neuroblastoma cell) cell lysate, COS-1 (African green monkey kidney fibroblast) cell lysate, PC-12 (Rat pheochromocytoma cell (undifferentiated)) cell lysate, Mouse brain tissue lysate, Rat brain tissue lysate, Pig brain tissue lysate.

Subcellular location: Mitochondrion, mitochondrion outer membrane, endoplasmic reticulum, cytosol, Nucleus, neuron projection, postsynaptic density, presynapse.

Database links: SwissProt: O60260 Human | Q9WS6 Mouse | Q9JK66 Rat

Recommended Dilutions:

WB	1:5,000
IF-Cell	1:100
IHC-P	1:2,000
IF-Tissue	1:500
FC(Intra)	1:1,000

Storage Buffer: 1*PBS (pH7.4), 0.1% BSA, 40% Glycerol, 0.2% Proclean 950.

Storage Instruction: Shipped at 4℃. Store at +4℃ short term (1-2 weeks). Store at -20℃ long term.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

Images

Fig1: Western blot analysis of Parkin on different lysates with Guinea pig anti-Parkin antibody (HA601663) at 1/5,000 dilution.



Lane 1: SH-SY5Y (Human neuroblastoma cell) cell lysate

Lane 2: COS-1 (African green monkey kidney fibroblast) cell lysate

Lane 3: PC-12 (Rat pheochromocytoma cell (undifferentiated)) cell lysate

Lane 4: Mouse brain tissue lysate

Lane 5: Rat brain tissue lysate

Lane 6: Pig brain tissue lysate

Lysates/proteins at 10 µg/Lane.

Exposure time: 10 seconds; ECL: K1801

Blocking: 5% NFDM/TBST, 1 hour at room temperature

Primary antibody: HA601663, 1/5,000 in primary antibody dilution buffer (K1803), overnight at 4 °C

Secondary antibody: Goat anti-Guinea pig IgG-HRP (HA1021), 1/50,000 in 5% NFDM/TBST, 1 hour at room temperature


Predicted band size: 52 kDa

Observed band size: 52 kDa

Fig2: Application: Immunocytochemistry (IF-cell)

Species: Mouse

Sample: Neuro-2a (Mouse brain neuroblastoma cell)



Fixation: 4% Paraformaldehyde, 15 minutes at room temperature.

Permeabilization: 0.1% Triton X-100, 15 minutes at room temperature.

Blocking: 1% BSA + 10% normal goat serum, 1 hour at room temperature.

Antibody dilution buffer: 1% BSA in PBST.

Primary antibody: HA601663, 1/100, overnight at 4°C.

Secondary antibody: Goat Anti-Guinea pig IgG (iFluor™ 594, HA1143), 45 hours at room temperature.

Counterstain: Beta tubulin (HA601187, Green), 1/100, overnight at 4°C. The nuclear counterstain was DAPI (Blue).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn




Fig3: Application: Immunohistochemistry (IHC-P)

Species: Mouse

Tissue: Cerebral cortex

Sample: Paraffin-embedded section

Antigen retrieval: Heat-mediated, Tris-EDTA buffer (pH 9.0), 20 minutes at 95°C.

Wash buffer: 1× TBST

Endogenous peroxidase blocking: 3% H₂O₂, 10 minutes at room temperature.

Blocking: 1% BSA + 10% normal goat serum, 10 minutes at room temperature.

Primary antibody: HA601663, 1/2,000, 1 hour at room temperature.

Secondary antibody: HA1021, 30 minutes at room temperature.




Fig4: Application: Immunohistochemistry (IHC-P)

Species: Rat

Tissue: Cerebral cortex

Sample: Paraffin-embedded section

Antigen retrieval: Heat-mediated, Tris-EDTA buffer (pH 9.0), 20 minutes at 95°C.


Wash buffer: 1× TBST

Endogenous peroxidase blocking: 3% H₂O₂, 10 minutes at room temperature.

Blocking: 1% BSA + 10% normal goat serum, 10 minutes at room temperature.

Primary antibody: HA601663, 1/2,000, 1 hour at room temperature.

Secondary antibody: HA1021, 30 minutes at room temperature.

**Fig5:** Application: Immunofluorescence (IF-tissue)

Species: Mouse

Tissue: Cerebral cortex

Sample: Paraffin-embedded section


Antigen retrieval: Heat-mediated, Tris-EDTA buffer (pH 9.0), 20 minutes at 95°C.

Wash buffer: 1× TBST

Blocking: 10% normal goat serum + 1% Triton X-100 + 0.3 M Glycine in TBST, 30 minutes at room temperature.

Primary antibody: HA601663, 1/500, overnight at 4°C.

Secondary antibody: Goat Anti-Guinea pig IgG (iFluor™ 594, HA1143), 1.5 hours at room temperature.

**Fig6:** Application: Flow Cytometry (Intra)

Species: Mouse

Sample: Neuro-2a (Mouse brain neuroblastoma cell)

Fixation: 4% Paraformaldehyde, 15 minutes at room temperature.

Permeabilization: 0.1% Tween-20, 15 minutes at room temperature.

Blocking: 1% BSA + 10% normal goat serum, 15 minutes at room temperature.

Antibody dilution buffer: 1x PBS.

Primary antibody: HA601663 (1/1,000, Red), 15 minutes at room temperature.

Secondary antibody: Goat Anti-Guinea pig IgG (iFluor™ 594, HA1143), 15 hours at room temperature.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Quinn PMJ et al. PINK1/PARKIN signalling in neurodegeneration and neuroinflammation. Acta Neuropathol Commun. 2020 Nov
2. Li J et al. PINK1/Parkin-mediated mitophagy in neurodegenerative diseases. Ageing Res Rev. 2023 Feb

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation