

Anti-T7 tag Antibody [4-C2-R] - BSA and Azide free

HA610073



Product Type:	Recombinant Mouse monoclonal IgG1, primary antibodies
Species reactivity:	Species independent
Applications:	WB
Clone number:	4-C2-R

Description: The T7 tag is an epitope tag composed of an 11-residue peptide encoded from the leader sequence of the T7 bacteriophage gene10. This gene encodes a T7 major capsid protein whose function is not clear. The T7 tag has been used extensively as a general epitope tag in many expression vectors including the pET system that is based on T7 RNA polymerase expression systems. This antibody recognizes the T7-tag fused to either the amino- or carboxy-terminus of targeted proteins.

Immunogen: Synthetic peptide CMASMTGGQMG.

Recommended Dilutions:
WB 1:1,000

Storage Buffer: 1*PBS (pH7.4).

Storage Instruction: Store at +4℃ after thawing. Aliquot store at -20℃. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

Images




Fig1: Western blot analysis of T7 tag on different T7 tagged recombinant proteins with Mouse anti-T7 tag antibody (HA610073) at 1/1,000 dilution.

Lysates/proteins at 50 ng/Lane.

Exposure time: 43 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA610073) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/50,000 dilution was used for 1 hour at room temperature.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Horn AV et al. Human LINE-1 restriction by APOBEC3C is deaminase independent and mediated by an ORF1p interaction that affects LINE reverse transcriptase activity. *Nucleic Acids Res* N/A:N/A (2013)
2. Kubota T., et al. Virus infection triggers SUMOylation of IRF3 and IRF7, leading to the negative regulation of type I interferon gene expression. *J Biol Chem.* 283:25660-70 (2008).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation