

Anti-VGLUT1 Antibody [PSH08-34] - BSA and Azide free

HA610217



Product Type: Recombinant Mouse monoclonal IgG1, primary antibodies

Species reactivity: Mouse, Rat

Applications: WB, IHC-P

Molecular Wt: Predicted band size: 62 kDa

Clone number: PSH08-34

Description: Vesicular glutamate transporter 1 (VGLUT1) is a protein that in humans is encoded by the SLC17A7 gene. The protein encoded by this gene is a vesicle-bound, sodium-dependent phosphate transporter that is specifically expressed in the neuron-rich regions of the brain. It is preferentially associated with the membranes of synaptic vesicles and functions in glutamate transport. The protein shares 82% identity with the differentiation-associated Na-dependent inorganic phosphate cotransporter and they appear to form a distinct class within the Na⁺/Pi cotransporter family.

Immunogen: Recombinant protein within rat VGlut1 aa 470-560.

Positive control: Mouse brain tissue lysate, Rat brain tissue lysate, mouse striatum tissue, mouse cerebellum tissue, rat striatum tissue, rat cerebellum tissue.

Subcellular location: Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane, Cell membrane, Synapse, synaptosome.

Database links: SwissProt: Q3TXX4 Mouse | Q62634 Rat

Recommended Dilutions:

WB 1:2,000

IHC-P 1:500

Storage Buffer: PBS (pH7.4).

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co.,Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images

Fig1: Western blot analysis of VGLUT1 on different lysates with Mouse anti-VGLUT1 antibody (HA610217) at 1/2,000 dilution.

Lane 1: Mouse brain tissue lysate (no heat) (20 µg/Lane)
 Lane 2: Rat brain tissue lysate (no heat) (20 µg/Lane)

Notice: no heat means the lysate is not boiled.

Predicted band size: 62 kDa
 Observed band size: 62 kDa

Exposure time: 14 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA610217) at 1/2,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/50,000 dilution was used for 1 hour at room temperature.




Fig2: Immunohistochemical analysis of paraffin-embedded mouse striatum tissue with Mouse anti-VGLUT1 antibody (HA610217) at 1/500 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA610217) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig3: Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue with Mouse anti-VGLUT1 antibody (HA610217) at 1/500 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA610217) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn




Fig4: Immunohistochemical analysis of paraffin-embedded rat striatum tissue with Mouse anti-VGLUT1 antibody (HA610217) at 1/500 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA610217) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig5: Immunohistochemical analysis of paraffin-embedded rat cerebellum tissue with Mouse anti-VGLUT1 antibody (HA610217) at 1/500 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA610217) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Souter EA et al. Disruption of VGLUT1 in Cholinergic Medial Habenula Projections Increases Nicotine Self-Administration. *eNeuro*. 2022 Jan
2. Jin S et al. Molecular Profiling of VGlut1 AND VGlut2 Ventral Subiculum to Nucleus Accumbens Shell Projections. *Neurochem Res*. 2023 Aug

Hangzhou Huan Biotechnology Co.,Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn