

HA610265



Product Type:	Recombinant Chimeric Antibody, primary antibodies
Species reactivity:	Mouse, Rat, Human, Cynomolgus monkey, Pig
Applications:	IHC-Fr, IF-Tissue
Molecular Wt:	Predicted band size: 13 kDa
Clone number:	PSH08-24

Description: Oxytocin is a peptide hormone and neuropeptide normally produced in the hypothalamus and released by the posterior pituitary. Present in animals since early stages of evolution, in humans it plays roles in behavior that include social bonding, love, reproduction, childbirth, and the period after childbirth. Oxytocin is released into the bloodstream as a hormone in response to sexual activity and during childbirth. It is also available in pharmaceutical form. In either form, oxytocin stimulates uterine contractions to speed up the process of childbirth. In its natural form, it also plays a role in maternal bonding and milk production. Production and secretion of oxytocin is controlled by a positive feedback mechanism, where its initial release stimulates production and release of further oxytocin. For example, when oxytocin is released during a contraction of the uterus at the start of childbirth, this stimulates production and release of more oxytocin and an increase in the intensity and frequency of contractions. This process compounds in intensity and frequency and continues until the triggering activity ceases. A similar process takes place during lactation and during sexual activity.

Immunogen:	Recombinant protein within human OXT aa 1-125.
Positive control:	Mouse paraventricular nucleus tissue, mouse pituitary tissue, rat pituitary tissue.
Subcellular location:	Secreted.
Database links:	SwissProt: P35454 Mouse Entrez Gene: 25504 Rat

Recommended Dilutions:

IHC-Fr	1:500
IF-Tissue	1:500-1:2,000

Storage Buffer: 1*PBS (pH7.4).

Storage Instruction: Store at +4℃ after thawing. Aliquot store at -20℃. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images

**Fig1:** Application: IHC-Fr


Species: Mouse

Site: Paraventricular nucleus

Sample: Frozen section

Antibody concentration: 1/500

Antigen retrieval: Not required


**Fig2:** Application: IF-tissue

Species: Mouse

Site: Paraventricular nucleus

Sample: Paraffin-embedded section

Antibody concentration: 1/500


**Fig3:** Application: IF-tissue

Species: Mouse

Site: Pituitary

Sample: Paraffin-embedded section

Antibody concentration: 1/500

**Fig4:** Application: IF-tissue

Species: Rat

Site: Pituitary

Sample: Paraffin-embedded section

Antibody concentration: 1/500

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. De Leon D et al. Methylation of OXT and OXTR genes, central oxytocin, and social behavior in female macaques. *Horm Behav.* 2020 Nov
2. Khazaal NM et al. The relationship between OXT gene polymorphisms and reproductive hormones in pregnant and lactating Awassi Ewes. *Mol Biol Rep.* 2023 Oct

Hangzhou Huaan Biotechnology Co., Ltd.

Orders: 0086-571-88062880

Technical: 0086-571-89986345

Service mail: support@huabio.cn

 华安生物
HUAABIO
www.huabio.cn

Applications: WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation