

HA610439



Product Type:	Recombinant Chimeric Antibody, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IHC-P
Molecular Wt:	Predicted band size: 12 kDa
Clone number:	JM100-08

Description: Parvalbumin (PV) is a calcium-binding protein with low molecular weight (typically 9-11 kDa). In humans, it is encoded by the PVALB gene. It is not a member of the albumin family; it is named for its size (parv-, from Latin parvus small) and its ability to coagulate. It has three EF hand motifs and is structurally related to calmodulin and troponin C. Parvalbumin is found in fast-contracting muscles, where its levels are highest, as well as in the brain and some endocrine tissues. Parvalbumin is a small, stable protein containing EF-hand type calcium binding sites. It is involved in calcium signaling. Calcium binding proteins like parvalbumin play a role in many physiological processes, namely cell-cycle regulation, second messenger production, muscle contraction, organization of microtubules and phototransduction. Therefore, calcium-binding proteins must distinguish calcium in the presence of high concentrations of other metal ions. The mechanism for the calcium selectivity has been extensively studied.

Immunogen: Recombinant protein within Human Parvalbumin aa 1-110 / 110.

Positive control: RPMI8226 (Human multiple myeloma cell) cell lysate, Mouse skeletal muscle tissue lysate, rat skeletal muscle tissue lysates.

Subcellular location: Axon, cytoplasm, synapse.

Database links: SwissProt: P20472 Human | P32848 Mouse | P02625 Rat

Recommended Dilutions:

WB	1:5,000
IHC-P	1:5,000

Storage Buffer: PBS (pH7.4).

Storage Instruction: Store at +4℃ after thawing. Aliquot store at -20℃. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images

Fig1: Western blot analysis of Parvalbumin on different lysates with Guinea pig anti-Parvalbumin antibody (HA610439) at 1/5,000 dilution.

Lane 1: RPMI8226 (Human multiple myeloma cell) cell lysate
Lane 2: SK-MEL-28 (Human malignant melanoma cell) cell lysate
Lane 3: Mouse skeletal muscle tissue lysate



Lysates/proteins at 15 µg/Lane.

Exposure time: 10 seconds; ECL: K1801

Negative expression of Parvalbumin protein in SK-MEL-28 cell is consistent with the predicted expression pattern.

Blocking: 5% NFDM/TBST, 1 hour at room temperature


Primary antibody: HA610439, 1/5,000 in primary antibody dilution buffer (K1803), overnight at 4 °C

Secondary antibody: Rabbit anti-Guinea pig IgG-HRP (HA1021), 1/50,000 in 5% NFDM/TBST, 1 hour at room temperature

Predicted band size: 12 kDa

Observed band size: 12 kDa

Fig2: Western blot analysis of Parvalbumin on rat skeletal muscle tissue lysate with Guinea pig anti-Parvalbumin antibody (HA610439) at 1/5,000 dilution.



Lysates/proteins at 15 µg/Lane.

Exposure time: 10 seconds; ECL: K1801

Blocking: 5% NFDM/TBST, 1 hour at room temperature

Primary antibody: HA610439, 1/5,000 in primary antibody dilution buffer (K1803), overnight at 4 °C

Secondary antibody: Rabbit anti-Guinea pig IgG-HRP (HA1021), 1/50,000 in 5% NFDM/TBST, 1 hour at room temperature

Predicted band size: 12 kDa

Observed band size: 12 kDa

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn




Fig3: Application: Immunohistochemistry (IHC-P)

Species: Mouse

Tissue: Cerebellum

Sample: Paraffin-embedded section

Antigen retrieval: Heat-mediated, Tris-EDTA buffer (pH 9.0), 20 minutes at 95°C.

Wash buffer: 1× TBST

Endogenous peroxidase blocking: 3% H₂O₂, 10 minutes at room temperature.

Blocking: 1% BSA + 10% normal goat serum, 10 minutes at room temperature.

Primary antibody: HA610439, 1/5,000, 1 hour at room temperature.

Secondary antibody: HA1021, 20 minutes at room temperature.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Hijazi S et al. Fast-spiking parvalbumin-positive interneurons in brain physiology and Alzheimer's disease. Mol Psychiatry. 2023 Dec
2. Feng YF et al. Parvalbumin neurons mediate neurological phenotypes of anti-NMDAR encephalitis. Brain. 2025 May

Hangzhou Huaan Biotechnology Co., Ltd.

Orders: 0086-571-88062880

Technical: 0086-571-89986345

Service mail: support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Applications: WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation